(19) Europäisches Patentamt
European Patent Office
Office européen des brevets

(11) EP 2 900 817 B1

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention of the grant of the patent:
10.11.2021 Bulletin 2021/45

(21) Application number: **13788679.2**

(22) Date of filing: **25.09.2013**

(51) Int Cl.:
A01H 1/06 *(2006.01)* ***A01H 5/08*** *(2018.01)*
C12N 9/88 *(2006.01)*

(86) International application number:
PCT/EP2013/069985

(87) International publication number:
WO 2014/049002 (03.04.2014 Gazette 2014/14)

(54) **SOLANUM LYCOPERSICUM PLANTS HAVING NON-TRANSGENIC ALTERATIONS IN THE ACS4 GENE**

SOLANUM LYCOPERSICUM-PFLANZEN MIT NICHT-TRANSGENEN VERÄNDERUNGEN IM ACS4-GEN

PLANTS DE SOLANUM LYCOPERSICUM PRÉSENTANT DES MODIFICATIONS NON TRANSGÉNIQUES DANS LE GÈNE ACS4

(84) Designated Contracting States:
AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR

(30) Priority: **28.09.2012 EP 12186606**

(43) Date of publication of application:
05.08.2015 Bulletin 2015/32

(83) **Declaration under Rule 32(1) EPC (expert solution)**

(73) Proprietor: **Nunhems B.V.**
6083 AB Nunhem (NL)

(72) Inventor: **VRIEZEN, Willem Hendrik**
NL-6081 BM Haelen (NL)

(74) Representative: **BASF IP Association**
BASF SE
G-FLP-C006
67056 Ludwigshafen (DE)

(56) References cited:
WO-A1-97/01952
WO-A2-2005/016504
WO-A2-2007/044043
US-A- 5 723 766

- **TSUCHISAKA ATSUNARI ET AL: "A Combinatorial Interplay Among the 1-Aminocyclopropane-1-Carboxylate Isoforms Regulates Ethylene Biosynthesis in Arabidopsis thaliana", GENETICS, vol. 183, no. 3, November 2009 (2009-11), pages 979-1003, XP002691844, ISSN: 0016-6731**
- **LINCOLN JAMES E ET AL: "LE-ACS4, a fruit ripening and wound-induced 1-aminocyclopropane-1-carboxylate synthase gene of tomato (Lycopersicon esculentum): Expression in Escherichia coli structural characterization, expression characteristics, and phylogenetic analysis", JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 268, no. 26, 1993, pages 19422-19430, XP002691845, ISSN: 0021-9258**
- **TARUN ALICE S ET AL: "Complementation analysis of mutants of 1-aminocyclopropane-1-carboxylate synthase reveals the enzyme is a dimer with shared active sites", JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 273, no. 20, 15 May 1998 (1998-05-15) , pages 12509-12514, XP002691846, ISSN: 0021-9258**
- **MINOIA SILVIA ET AL: "A new mutant genetic resource for tomato crop improvement by TILLING technology", BMC RESEARCH NOTES, BIOMED CENTRAL LTD, GB, vol. 3, no. 69, 1 January 2010 (2010-01-01), pages 1-8, XP002682656, ISSN: 1756-0500**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 2 900 817 B1

Description

FIELD OF THE INVENTION

[0001] This invention relates to the field of plant biotechnology and plant breeding. Provided are *Solanum lycopersicum* plants comprising an *acs4* allele having one or more mutations, said mutations resulting in production of a mutant acs4 protein having loss-of-function acs4 protein or reduced activity compared to wild type Acs4 protein. The invention provides plants the fruits of which show a lower ethylene production and/or slower fruit ripening and/or a longer shelf life compared to *Solanum lycopersicum* being homozygous for the wild type *Acs4* allele. In addition, the invention provides tomato fruit, seeds, pollen, plant parts, and progeny of the *Solanum lycopersicum* plants of the invention. Food and food products comprising or consisting of fruits of the plants of the invention are provided too.

[0002] The invention further provides an endogenous *acs4* gene and acs4 protein encoded by said gene, having at least one human-induced non-transgenic mutation.

[0003] In another embodiment methods for making tomato plants comprising one or more mutant *acs4* alleles in their genome are provided herein.

BACKGROUND OF THE INVENTION

[0004] Breeding of *Solanum lycopersicum* aims at the production of commercial varieties optimally adapted to growing and storage conditions. A challenge breeders are facing is finding an improved balance between fruit firmness post-harvest and consumer desires in terms of taste, texture and colour. These consumer desires relate strongly to fruit ripening. Fruit ripening is a complex developmental process responsible for the transformation of the seed-containing organ into a tissue attractive to seed dispersers and agricultural consumers. The changes associated with fruit ripening, in particular post-harvest softening, limit the shelf life of fresh tomatoes.

[0005] For tomato fruit growth and development, a number of consecutive phases can be discerned: floral development, pollination, then early fruit development takes place which is characterised by a high frequency of cell division and the fruit is rapidly increasing in size mainly due to cell expansion. At the end of the third phase the fruit reaches the mature green stage. During the fourth phase, fruit ripening takes place which is characterised by a change in colour and flavour as well as fruit firmness and texture.

[0006] The build-up of the characteristic red colour of the tomato fruit is caused by the accumulation of lycopene and carotene. In general, different colouration phases are distinguished: mature green, breaker, pink and red. At the breaker stage, the typical red pigmentation initiates. Red ripe stage or red ripe harvested fruit stage is the stage where the fruit has reached its mature colour on the major part of the fruit.

[0007] In addition to the colour changes, during fruit ripening enzymatic activity leads to degradation of the middle lamellar region of the cell walls which leads to cell loosening which is manifested as softening and loss of texture of the fruit. Softening of the fruit is often measured as external resistance to compression which can be quantified for example by a penetrometer.

[0008] Modification of single genes known to be involved in ripening has not yet resulted in a fruit with normal ripening but minimal tissue softening.

[0009] Ripening and senescence in climacteric fruits such as tomatoes are promoted by ethylene. Ethylene is auto-catalytic for its own biosynthesis through increases in 1-Aminocyclopropae-1-carboxilic acid (ACC) synthase (ACS) and ACC oxidase (ACO). ACS is also referred to as 1-aminocyclopropane-1-carboxylate synthase; Le-ACS; or S-adenosyl-L-methionine methylthioadenosine-lyase. An increase in the amount of ACS and ACO thus leads to an increased conversion of L-methionine into ethylene. At least eight ACS genes (LEACS1A, LEACS1B, and LEACS2-7) have been identified in tomato (Alexander et. al., Journal of Experimental Botany, Vol 53, No 377, pp 2039-2055, 2002) and each ACS has a different expression pattern.

[0010] ACC synthase (ACS) is an enzyme that catalyzes the synthesis of 1-aminocyclopropane-1-carboxylic acid (ACC) from S-Adenosyl methionine. ACC is then converted into ethylene catalyzed by ACO. The biosynthesis of ethylene is for example described by Stearns and Glick (Biotechnology Advances 2003, vol 21 pp 193-210).

[0011] ACS belongs to the α-family of pyridoxal-5'-phosphate (PLP) dependent enzymes and shares a modest level of similarity with other members of this family like aspartate amino-transferase (AATase and tyrosine aminotransferase (TATase). The structure of ACS from various sources has been described by Capitani et al. In a sequence alignment of eight ACS proteins (Malus domestica, Phaseolus aureus, Solanum tuberosum, Pelargonium hortorum, Nicotiana tabacum, Cucumis melo, Lycopersicon esculentum, and Brassica oleracea*)* they describe conserved regions which are indicated in red and yellow in Figure 1 in this Capitani publication. Three domains are defined: one large domain ranging from residue 52 to 318 and two small domains, ranging from residues 20 to 49 and 333 to 430. An helix α12 is defined connecting the large domain with the second small domain (Capitani et al., Journal of Molecular Biology, 1999, vol 294, pp 745-756).

[0012] Two systems have been proposed to operate in climacteric plants regulating ethylene production. The first is functional during normal vegetative growth (system 1); it is auto inhibitory and responsible for production of basal ethylene levels that are detected in all tissues including those in non-climacteric plants. System 1 continues during fruit development until a competence to fruit ripening is attained. Then a transition period is reached wherein LEACS1A and LEACS4 are activated resulting in an increased level of ethylene. This increased ethylene level induces the expression of LEACS2 which starts system 2 which is active during the ripening of climacteric fruit. In system 2, ethylene production is auto catalytic. This complexity of the ethylene regulation has been studied using antisense inhibition of LEACS2 in transgenic plants (Barry et al., Plant Physiology vol 123, pp 979-986, 2000).

[0013] WO2005/016504 discloses "stay green" plants, i.e. a plant phenotype whereby leaf senescence is delayed compared to a standard reference. It discloses plants with disrupted ACS2, ACS6, ASC7 genes which disruption inhibits the expression or activity of said ACS.

[0014] Yokotani *et al* describe transgenic tomatoes with all known LeEIL genes (Ethylene Insensitive Like genes) suppressed to study the regulatory mechanisms of ethylene biosynthesis (Yokotani et al, Journal of Experimental Botany, vol 60, pp 3433-3442, 2009).

[0015] There is thus a need for cultivated tomato plants with a modified ethylene production having a delayed ripening and/or longer shelf-life of the tomato fruits compared to wild type tomato plants.

SUMMARY OF THE INVENTION

[0016] It is, thus, an object of the invention to generate and identify cultivated plants of the species *Solanum lycopersicum* having fruits that have delayed ripening and/or a longer shelf-life of the fruits.

[0017] The invention thus relates to a cultivated plant of the species *Solanum lycopersicum* comprising an *acs4* allele having one or more mutations, said mutations resulting in production of a mutant acs4 protein having loss-of-function acs4 protein or reduced activity compared to wild type Acs4 protein of SEQ ID NO: 1 but which comprises sufficient function to result in ripening of the tomato fruits to the red stage when the mutant allele is present in heterozygous or homozygous form.

GENERAL DEFINITIONS

[0018] The term "nucleic acid sequence" (or nucleic acid molecule) refers to a DNA or RNA molecule in single or double stranded form, particularly a DNA encoding a protein or protein fragment according to the invention. An "isolated nucleic acid sequence" refers to a nucleic acid sequence which is no longer in the natural environment from which it was isolated, e.g. the nucleic acid sequence in a bacterial host cell or in the plant nuclear or plastid genome.

[0019] The terms "protein" or "polypeptide" are used interchangeably and refer to molecules consisting of a chain of amino acids, without reference to a specific mode of action, size, 3-dimensional structure or origin. A "fragment" or "portion" of Acs4 protein may thus still be referred to as a "protein". An "isolated protein" is used to refer to a protein which is no longer in its natural environment, for example *in vitro* or in a recombinant bacterial or plant host cell.

[0020] The term "gene" means a DNA sequence comprising a region (transcribed region), which is transcribed into an RNA molecule (e.g. an mRNA or an RNAi molecule) in a cell, operably linked to suitable regulatory regions (e.g. a promoter). A gene may thus comprise several operably linked sequences, such as a promoter, a 5' leader sequence comprising e.g. sequences involved in translation initiation, a (protein) coding region (cDNA or genomic DNA) and a 3' non-translated sequence comprising e.g. transcription termination sites. A gene may be an endogenous gene (in the species of origin) or a chimeric gene (e.g. a transgene or cis-gene).

[0021] "Expression of a gene" refers to the process wherein a DNA region, which is operably linked to appropriate regulatory regions, particularly a promoter, is transcribed into an RNA, which is biologically active, i.e. which is capable of being translated into a biologically active protein or peptide (or active peptide fragment) or which is active itself (e.g. in posttranscriptional gene silencing or RNAi). The coding sequence may be in sense-orientation and encodes a desired, biologically active protein or peptide, or an active peptide fragment.

[0022] An "active protein" or "functional protein" is a protein which has protein activity as measurable *in vitro,* e.g. by an *in vitro* activity assay, and/or *in vivo,* e.g. by the phenotype conferred by the protein. A "wild type" protein is a fully functional protein, as present in the wild type plant. A "mutant protein" is herein a protein comprising one or more mutations in the nucleic acid sequence encoding the protein, whereby the mutation results in (the mutant nucleic acid molecule encoding) a "reduced-function" or "loss-of-function" protein, as e.g. measurable *in vivo,* e.g. by the phenotype conferred by the mutant allele.

[0023] A "reduced function acs4 protein" or "reduced activity acs4 protein" refers to a mutant acs4 protein which has a reduced catalytic activity in synthesizing ACC from S-Adenosyl methionine, leading to reduced ethylene synthesis compared to wild-type Acs4 protein. Said reduced catalytic activity of the acs4 protein affects the ripening behaviour of the fruits comprising such reduced function acs4 protein when the allele encoding the mutant protein is present in

homozygous or heterozygous form in the tomato plant, i.e. delayed ripening and/or longer shelf-life of the fruits. Such a reduced function acs4 protein can be obtained by the transcription and translation of a "partial knockout mutant *acs4* allele" which is, for example, a wild-type *Acs4* allele, which comprises one or more mutations in its nucleic acid sequence. In one aspect, such a partial knockout mutant *acs4* allele is a wild-type *Acs4* allele, which comprises one or more mutations that preferably result in the production of an acs4 protein wherein at least one conserved and/or functional amino acid is substituted for another amino acid, such that the biological activity is significantly reduced but not completely abolished. However, other mutations, such as one or more non-sense, missense, splice-site or frameshift mutations in the tomato *Acs4* allele may also result in reduced function acs4 protein and such reduced function proteins may have one or more amino acids replaced, inserted or deleted, relative to the wild type ACS4 protein. Such partial knockout mutant *acs4* allele may also encode a dominant negative acs4 protein, which is capable of adversely affecting the biological activity of other Acs4 proteins within the same cell. Such a dominant negative acs4 protein can be an acs4 protein that is still capable of interacting with the same elements as the wild-type Acs4 protein, but that blocks some aspect of its function. Examples of dominant negative acs4 proteins are acs4 proteins that lack, or have modifications in specific amino acid residues critical for activation, but still contain their binding domain, such that not only their own biological activity is reduced or abolished, but that they further reduce the total acs4 activity in the cell by competing with wild type and/or partial knockout acs4 proteins present in the cell for binding sites. Mutant alleles can be either "natural mutant" alleles, which are mutant alleles found in nature (e.g. produced spontaneously without human application of mutagens) or "induced mutant" alleles, which are induced by human intervention, e.g. by mutagenesis.

[0024] A "loss-of-function acs4 protein" refers to a mutant acs4 protein which has essentially no catalytic activity in synthesising ACC from S-Adenosyl methionine compared to wild-type Acs4 protein, leading to reduced ethylene synthesis compared to wild type Acs4 protein. Said lack of catalytic activity synthesis affects the ripening behaviour of the fruits comprising such loss-of-function acs4 protein when the allele encoding the mutant protein is present in homozygous or heterozygous form in the tomato plant. Fruits of tomato plants homozygous for such a "loss-of- function acs4 protein" may still produce ethylene catalysed by other proteins (e.g. other Acs proteins like Acs1A). As a consequence, fruits of tomato plants homozygous for such a "loss-of-function acs4 protein" may still ripen, but ripening may be delayed and/or shelf life may be longer.

[0025] A "mutation" in a nucleic acid molecule coding for a protein is a change of one or more nucleotides compared to the wild type sequence, e.g. by replacement, deletion or insertion of one or more nucleotides. A "point mutation" is the replacement of a single nucleotide, or the insertion or deletion of a single nucleotide.

[0026] A "nonsense" mutation is a (point) mutation in a nucleic acid sequence encoding a protein, whereby a codon is changed into a stop codon. This results in a premature stop codon being present in the mRNA and in a truncated protein. A truncated protein may have reduced function or loss of function.

[0027] A "missense" or non-synonymous mutation is a (point) mutation in a nucleic acid sequence encoding a protein, whereby a codon is changed to code for a different amino acid. The resulting protein may have reduced function or loss of function.

[0028] A "splice-site" mutation is a mutation in a nucleic acid sequence encoding a protein, whereby RNA splicing of the pre-mRNA is changed, resulting in an mRNA having a different nucleotide sequence and a protein having a different amino acid sequence than the wild type. The resulting protein may have reduced function or loss of function.

[0029] A "frame-shift" mutation is a mutation in a nucleic acid sequence encoding a protein by which the reading frame of the mRNA is changed, resulting in a different amino acid sequence. The resulting protein may have reduced function or loss of function.

[0030] A mutation in a regulatory sequence, e.g. in a promoter of a gene, is a change of one or more nucleotides compared to the wild type sequence, e.g. by replacement, deletion or insertion of one or more nucleotides, leading for example to reduced or no mRNA transcript of the gene being made.

[0031] "Silencing" refers to a down-regulation or complete inhibition of gene expression of the target gene or gene family.

[0032] A "target gene" in gene silencing approaches is the gene or gene family (or one or more specific alleles of the gene) of which the endogenous gene expression is down-regulated or completely inhibited (silenced) when a chimeric silencing gene (or 'chimeric RNAi gene') is expressed and for example produces a silencing RNA transcript (e.g. a dsRNA or hairpin RNA capable of silencing the endogenous target gene expression). In mutagenesis approaches, a target gene is the endogenous gene which is to be mutated, leading to a change in (reduction or loss of) gene expression or a change in (reduction or loss of) function of the encoded protein.

[0033] As used herein, the term "operably linked" refers to a linkage of polynucleotide elements in a functional relationship. A nucleic acid is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For instance, a promoter, or rather a transcription regulatory sequence, is operably linked to a coding sequence if it affects the transcription of the coding sequence. Operably linked means that the DNA sequences being linked are typically contiguous and, where necessary to join two protein encoding regions, contiguous and in reading frame so as to produce a "chimeric protein". A "chimeric protein" or "hybrid protein" is a protein composed of various protein "domains" (or motifs) which is not found as such in nature but which a joined to form a functional protein, which displays the

functionality of the joined domains. A chimeric protein may also be a fusion protein of two or more proteins occurring in nature.

[0034] The term "food" is any substance consumed to provide nutritional support for the body. It is usually of plant or animal origin, and contains essential nutrients, such as carbohydrates, fats, proteins, vitamins, or minerals. The substance is ingested by an organism and assimilated by the organism's cells in an effort to produce energy, maintain life, or stimulate growth. The term food includes both substance consumed to provide nutritional support for the human and animal body.

[0035] The term "shelf life" or "post-harvest shelf life" designates the (average) length of time that a fruit is given before it is considered unsuitable for sale or consumption ('bad'). Shelf life is the period of time that products can be stored, during which the defined quality of a specified proportion of the goods remains acceptable under expected conditions of distribution, storage and display. Shelf life is influenced by several factors: exposure to light and heat, transmission of gases (including humidity), mechanical stresses, and contamination by things such as micro-organisms. Product quality is often mathematically modelled around the fruit firmness/softness parameter. Shelf-life can be defined as the (average) time it takes for fruits of a plant line to start to become bad and unsuitable for sale or consumption, starting for example from the first fruit of a plant entering breaker stage or turning stage or from the first fruit becoming fully red or from harvest. In one embodiment the mutants according to the invention have a shelf life that is significantly longer than the shelf life of wild type plants, for example the number of days from the first fruit being in breaker stage (or turning stage, pink stage, red stage or from harvest) up to the first fruit starting to become 'bad' and unsuitable for sale or consumption is significantly longer, e.g. at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more, days longer than fruits of control plants (such as wild type *Acs4*/*Acs4* plants), when plants are grown under the same conditions and fruits are treated the same way and kept under the same conditions. Thus, to determine the number of days required from a certain stage (e.g. from breaker stage or a later stage) to 'bad' stage, the day when the first fruit of the wild type control plant (grown under the same conditions as the mutant plants and being at the same developmental stage) enters a certain stage (e.g. breaker stage or a later stage) can, for example, be taken as the starting point (day 1) from when on periodically (at certain time intervals, e.g. after 1, 2, 3, 4, 5 or 6 days) the fruits are observed until the day that the first fruit has passed the fully ripe stage and becomes 'bad' (as determinable visually and/or through assessing fruit softness).

[0036] In this application the words "improved", "increased", "longer" and "extended" as used in conjunction with the word "shelf-life" are interchangeable and all mean that the fruits of a tomato plant according to the invention have on average, a longer shelf-life than the control fruits *(Acs4*/*Acs4* fruits).

[0037] "Delayed ripening" means that the fruits of a tomato plant or plant line (e.g. a mutant) according to the invention require on average significantly more days to reach the red stage from the mature green, breaker, turning stage, and/or pink stages of tomato fruit ripening compared to wild type control fruits of plants homozygous for the wild type *Acs4* allele *(Acs4*/*Acs4).* Delayed ripening can be measured on the plant and/or after harvest as days required for a certain percentage of fruits (e.g. 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% and/or 100% of fruits) to reach the red stage. A plant is said to have a delayed ripening phenotype if it takes at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15 days longer for 10%, 20%, 30% 40%, 50%, 60%, 70%, 80%, 90% and/or 100% of fruits to reach the red stage than it takes for the wild type control fruits to develop the same percentage of red fruits. It is understood that each combination of above-cited number of days (i.e. 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15) with each % of fruits to reach the red stage (i.e. 10%, 20%, 30% 40%, 50%, 60%, 70%, 80%, 90% and/or 100%) is enclosed herein, both for the delayed ripening to be measured on the plant and after harvest. For example if it takes at least 2 days longer for 10%, 20%, 30% 40%, 50%, 60%, 70%, 80%, 90% and/or 100% of fruits to reach the red stage than it takes for the wild type control fruits to develop the same percentage of red fruits. Another example of how delayed ripening can be measured on the plant and/or after harvest is it takes at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15 days longer for 100% of fruits to reach the red stage than it takes for the wild type control fruits to develop the same percentage of red fruits. The day when the first fruit of the wild type control plant (grown under the same conditions as the mutant plants and being at the same developmental stage) enters a certain stage (e.g. breaker stage) can, for example, be taken as the starting point (day 1) from when on periodically (at certain time intervals (e.g. after 1, 2, 3, 4, 5 or 6 days) the number of fruits that are in breaker stage and the number of fruit that are in red stage are counted, both for the mutant plant line and control plants (see Examples).

[0038] As used herein, "reduced ethylene production" refers herein to statistically significant reduced amounts of ethylene being produced by tomato fruits according to the invention (compared to wild type Acs4/Acs4 fruits) during fruit ripening, e.g. at the pink stage and/or at the light red stage and/or at the red stage, as described in the Examples, and as measurable by real time ethylene measurements. In one embodiment, ethylene levels are significantly reduced throughout fruit ripening from pink stage through to red stage.

[0039] It is understood that comparisons between different plant lines involves growing a number of plants of a line (e.g. at least 5 plants, preferably at least 10 plants per line) under the same conditions as the plants of one or more control plant lines (preferably wild type plants) and the determination of statistically significant differences between the plant lines when grown under the same environmental conditions.

[0040] "Delay of breaker stage" refers to the mutants according to the invention requiring significantly more days than wild type controls for the first fruits and/or for all fruits to have entered breaker stage, e.g. at least 1 more day, preferably at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 more days than the wild type control, when grown under the same conditions.

[0041] The "ripening stage" of a tomato fruit can be divided as follows: (1) Mature green stage: surface is completely green; the shade of green may vary from light to dark. (2) Breaker stage: there is a definite break in color from green to tannish-yellow, pink or red on not more than 10% of the surface; (3) Turning stage: 10% to 30% of the surface is not green; in the aggregate, shows a definite change from green to tannish-yellow, pink, red, or a combination thereof. (4) Pink stage: 30% to 60% of the surface is not green; in the aggregate, shows pink or red color. (5) Light red stage: 60% to 90% of the surface is not green; in the aggregate, shows pinkish-red or red. (6) Red stage: More than 90% of the surface is not green; in the aggregate, shows red color.

[0042] "Sequence identity" and "sequence similarity" can be determined by alignment of two peptide or two nucleotide sequences using global or local alignment algorithms. Sequences may then be referred to as "substantially identical" or "essentially similar" when they are optimally aligned by for example the programs GAP or BESTFIT or the Emboss program "Needle" (using default parameters, see below) share at least a certain minimal percentage of sequence identity (as defined further below). These programs use the Needleman and Wunsch global alignment algorithm to align two sequences over their entire length, maximizing the number of matches and minimises the number of gaps. Generally, the default parameters are used, with a gap creation penalty = 10 and gap extension penalty = 0.5 (both for nucleotide and protein alignments). For nucleotides the default scoring matrix used is DNAFULL and for proteins the default scoring matrix is Blosum62 (Henikoff & Henikoff, 1992, PNAS 89, 10915-10919). Sequence alignments and scores for percentage sequence identity may for example be determined using computer programs, such as EMBOSS (http://www.ebi.ac.uk/Tools/psa/emboss_needle/). Alternatively sequence similarity or identity may be determined by searching against databases such as FASTA, BLAST, etc., but hits should be retrieved and aligned pairwise to compare sequence identity. Two proteins or two protein domains, or two nucleic acid sequences have "substantial sequence identity" if the percentage sequence identity is at least 90%, 95%, 98%, 99% or more (as determined by Emboss "needle" using default parameters, i.e. gap creation penalty = 10, gap extension penalty = 0.5, using scoring matrix DNAFULL for nucleic acids an Blosum62 for proteins). Such sequences are also referred to as 'variants' herein, e.g. other variants of mutant *acs4* alleles and mutant acs4 proteins than the specific nucleic acid and protein sequences disclosed herein can be identified, which have the same effect on delayed ripening and/or longer shelf-life of the fruits comprising such variants.

[0043] The amino acid sequence alignment of five of the sequences given in Figure 1 of Capitani et al. (Journal of Molecular Biology, 1999, vol 194, pp 745-756) (*Cucumis melo Accession Q42668, Pelargonium hortorum Accession Q43810, Brassica oleracea Accession Q43747, Phaseolus aureus Accession Q41688,* and *Solanum tuberosum Accession Q43166*) with the wild type *Solanum lycopersicum* ACS4 amino acid sequence as given in SEQ ID NO 1 (Le-ACS4) is shown in Figure 1 of this application. This alignment reveals, see Figure 1, that the conserved amino acids as indicated in yellow and red in Figure 1 of Capitani *et al.* are also conserved in wild type *Solanum lycopersicum* ACS4 amino acid sequence. Note that the amino acid numbering in Figure 1 of this application as indicated does not correspond to the numbering in Figure 1 of Capitani *et al.*.

[0044] The ACS4 "large domain" refers to amino acid residues from amino acid 65 to amino acid 327 of SEQ ID NO: 1 (see also Figure 4). The ACS4 small domains refer to either amino acid residues 33 to 62 of SEQ ID NO: 1 (see Figure 4) and/or from amino acid 339 to amino acid 438 of SEQ ID NO: 1 (see Figure 4) of this application. The ACS4 catalytic centre is believed to be in the "large domain".

[0045] In this document and in its claims, the verb "to comprise" and its conjugations is used in its non-limiting sense to mean that items following the word are included, but items not specifically mentioned are not excluded. In addition, reference to an element by the indefinite article "a" or "an" does not exclude the possibility that more than one of the element is present, unless the context clearly requires that there be one and only one of the elements. The indefinite article "a" or "an" thus usually means "at least one". It is further understood that, when referring to "sequences" herein, generally the actual physical molecules with a certain sequence of subunits (e.g. amino acids) are referred to.

[0046] As used herein, the term "plant" includes the whole plant or any parts or derivatives thereof, such as plant organs (e.g., harvested or non-harvested fruits, flowers, leaves, etc.), plant cells, plant protoplasts, plant cell or tissue cultures from which whole plants can be regenerated, regenerable or non-regenerable plant cells, plant calli, plant cell clumps, and plant cells that are intact in plants, or parts of plants, such as embryos, pollen, ovules, ovaries, fruits (e.g., harvested tissues or organs, such as harvested tomatoes or parts thereof), flowers, leaves, seeds, tubers, clonally propagated plants, roots, stems, cotyledons, hypocotyls, root tips and the like. Also any developmental stage is included, such as seedlings, immature and mature, etc.

[0047] A "plant line" or "breeding line" refers to a plant and its progeny. As used herein, the term "inbred line" refers to a plant line which has been repeatedly selfed.

[0048] "Plant variety" is a group of plants within the same botanical taxon of the lowest grade known, which (irrespective of whether the conditions for the recognition of plant breeder's rights are fulfilled or not) can be defined on the basis of

the expression of characteristics that result from a certain genotype or a combination of genotypes, can be distinguished from any other group of plants by the expression of at least one of those characteristics, and can be regarded as an entity, because it can be multiplied without any change. Therefore, the term "plant variety" cannot be used to denote a group of plants, even if they are of the same kind, if they are all characterized by the presence of 1 locus or gene (or a series of phenotypical characteristics due to this single locus or gene), but which can otherwise differ from one another enormously as regards the other loci or genes.

[0049] "F1, F2, etc." refers to the consecutive related generations following a cross between two parent plants or parent lines. The plants grown from the seeds produced by crossing two plants or lines is called the F1 generation. Selfing the F1 plants results in the F2 generation, etc. "F1 hybrid" plant (or F1 seed) is the generation obtained from crossing two inbred parent lines. An "M1 population" is a plurality of mutagenized seeds / plants of a certain plant line or cultivar. "M2, M3, M4, etc." refers to the consecutive generations obtained following selfing of a first mutagenized seed / plant (M1).

[0050] The term "allele(s)" means any of one or more alternative forms of a gene at a particular locus, all of which alleles relate to one trait or characteristic at a specific locus. In a diploid cell of an organism, alleles of a given gene are located at a specific location, or locus (loci plural) on a chromosome. One allele is present on each chromosome of the pair of homologous chromosomes. A diploid plant species may comprise a large number of different alleles at a particular locus. These may be identical alleles of the gene (homozygous) or two different alleles (heterozygous).

[0051] The term "locus" (loci plural) means a specific place or places or a site on a chromosome where for example a gene or genetic marker is found. The *ACS4* locus is thus the location in the genome where the *ACS4* gene is found.

[0052] "Wild type allele" (WT) refers herein to a version of a gene encoding a fully functional protein (wild type protein). Such a sequence encoding a fully functional Acs4 protein is for example the wild type Acs4 cDNA (mRNA) sequence depicted in SEQ ID NO: 8, based on GenBank Accession M63490.1 or the wild type *Acs4* genomic sequence depicted in SEQ ID NO: 15. The protein sequence encoded by this wild type *Acs4* mRNA is depicted in SEQ ID NO: 1 and in SEQ ID NO: 15. It consists of 476 amino acids. Three domains have been mentioned to occur on the Acs4 protein i.e. a first small domain ranging from amino acid 33 to 62 of SEQ ID NO: 1, a "large domain", presumed to contain the catalytic centre of the protein (ranging from amino acid 65 to 327 of SEQ ID NO: 1 and a second small domain ranging from amino acid residue 339 to 438of SEQ ID NO: 1 (see Figure 4). Other fully functional Acs4 protein encoding alleles (i.e. alleles which confer ripening and ethylene production to the same extent as the protein of SEQ ID NO 1) may exist in other *Solanum lycopersicum* plants and may comprise substantial sequence identity with SEQ ID NO: 1, i.e. at least about 90%, 95%, 98%, 99%, 99.2%, 99.3%, 99.4%, 99.5%, 99.6%, 99.7% sequence identity with SEQ ID NO: 1. Such fully functional wild type Acs4 proteins are herein referred to as "variants" of SEQ ID NO: 1. Likewise the nucleotide sequences encoding such fully functional Acs4 proteins are referred to as variants of SEQ ID NO: 8 and SEQ ID NO: 15.

[0053] The following mutant *acs4* alleles are exemplary of the reduced ethylene production and/or delayed-ripening and/or extended shelf-life conferring acs4 mutations identified according to the present invention. It is noted that nucleotide sequences referred to herein (SEQ ID NO: 8-14) are cDNA, i.e. coding DNA sequences, encoding the proteins of SEQ ID NO: 1-7. Obviously, when reference is made to these cDNA nucleotide sequences, it is understood that the cDNA is the coding region of the corresponding *Solanum lycopersicum* genomic *acs4* sequence, which, however, additionally contains introns and therefore the nucleotides have different numbering. Thus, when reference is made to a tomato plant comprising an *acs4* sequence according to e.g. any one of SEQ ID NO: 8-14, it is, therefore, understood that the tomato plant comprising the genomic *acs4* sequence which comprises the coding DNA (cDNA), from which the mRNA of SEQ ID NO: 8-14 is transcribed (and which is in turn translated into protein). The mRNA has the same nucleotide sequence as the cDNA, accept that Thymine (t) is Uracil (u) in the mRNA. Further, when reference is made to a tomato plant comprising a nucleotide sequence encoding a protein according to the invention (such as a mutant protein of SEQ ID No: 2-7, or a different mutant), this encompasses different nucleotide sequences, due to the degeneracy of the genetic code. In one embodiment the plant comprises the genomic Acs4 sequence depicted in SEQ ID NO: 15 or a genomic Acs4 sequence substantially identical thereto (e.g. having at least about 70%, 75%, 80%, 85%, 90%, 95%, 98%, 99%, 99.2%, 99.3%, 99.4%, 99.5%, 99.6%, 99.7% sequence identity with SEQ ID NO: 15), but with one or more mutations in said sequence, especially in the exons of said genomic sequence (exon 1 ranges from nucleotide 1 to 318; exon 2 ranges from nucleotide 796 to 955 and exon 3 ranges from nucleotide 1689 to 2638), causing reduced function or loss of function of the encoded mutant acs4 protein.

[0054] One exemplary mutant *acs4* allele (mutant 2477, or Nun 2477) conferring reduced ethylene production and/or delayed ripening and/or extended shelf-life identified according to the present invention, comprises a mutation resulting in a serine (Ser or S) to asparagine (Asn or N) substitution at amino acid 279 in the encoded protein (SEQ ID NO: 2). The S279N mutation is within the large-domain of the ACS4 protein. The protein sequence of mutant 2477 is depicted in SEQ ID NO: 2. The amino acid substitution is due to a G to A mutation at nucleotide 836 of SEQ ID NO: 8 counting A in the ATG of the START CODON as nucleotide position 1. The mutant cDNA is depicted in SEQ ID NO: 9.

[0055] Another exemplary mutant *acs4* allele (mutant 4043, or Nun 4043) conferring reduced ethylene production and/or delayed ripening and/or extended shelf-life identified according to the present invention, comprises a mutation

resulting in a change from alanine (Ala or A) to valine (Val or V) at amino acid 248 in the encoded protein (SEQ ID NO: 3). The A248V mutation is within the large-domain of the ACS4 protein. The protein sequence of mutant 4043 is depicted in SEQ ID NO: 3. The amino acid substitution is due to a C to T mutation at nucleotide 743 of SEQ ID NO: 1 , counting A in the ATG of the START CODON as nucleotide position 1. The mutant cDNA is depicted in SEQ ID NO: 10.

[0056] Still another exemplary mutant *acs4* allele (mutant 4222, or Nun 4222) conferring reduced ethylene production and /or delayed ripening and/or extended shelf-life, identified according to the present invention, comprises a mutation resulting in a truncated protein of 203 amino acid residues during translation, whereas the wild type protein has 476 amino acid residues. The truncated protein sequence of mutant 4222 is depicted in SEQ ID NO: 4. The truncation is due to a change from A to T at nucleotide 610 of SEQ ID NO: 1 counting A in the ATG of the START CODON as nucleotide position 1. This A610T mutation in mutant 4222 results in a change from a codon for lysine (AAA) to a STOP-codon (TAA). The mutant cDNA is depicted in SEQ ID NO: 11.

[0057] Another exemplary mutant *acs4* allele (mutant 4303 or Nun 4303) conferring reduced ethylene production and /or delayed ripening and/or extended shelf-life, identified according to the present invention, comprises a mutation resulting in a change from leucine (Leu or L) to phenylalanine (Phe or F) at amino acid 321 in the encoded protein. The L321F mutation is within the second small-domain of the ACS4 protein. The protein sequence of mutant 4303 is depicted in SEQ ID NO: 5. The amino acid substitution is due to a change from G to T at nucleotide 963 of SEQ ID NO: 1 counting A in the ATG of the START CODON as nucleotide position 1. The mutant cDNA is depicted in SEQ ID NO: 12.

[0058] Yet another exemplary mutant *acs4* allele (mutant 4691, or Nun 4691) conferring reduced ethylene production and /or delayed ripening and/or extended shelf-life, identified according to the present invention, comprises a mutation resulting in a change from valine (Val or V) to glutamic acid (Glu or E) at amino acid 250 in the encoded protein. The V250E mutation is within the large-domain of the ACS4 protein. The protein sequence of mutant 4691 is depicted in SEQ ID NO: 6. The amino acid substitution is due to a change from T to A at nucleotide 749 of as shown in SEQ ID NO: 1 counting A in the ATG of the START CODON as nucleotide position 1. The mutant cDNA is depicted in SEQ ID NO: 13.

[0059] Another exemplary mutant *acs4* allele (mutant 5251, or Nun 5251) conferring reduced ethylene production and /or delayed ripening and/or extended shelf-life, identified according to the present invention, comprises a mutation resulting in a change from threonine (Thr or T) to isoleucine (Ile or I) at amino acid 316 in the encoded protein. The T316I mutation is within the second small-domain of the ACS4 protein. The protein sequence of mutant 5251 is depicted in SEQ ID NO: 7. The amino acid substitution is due to a change from C to T at nucleotide 947 of SEQ ID NO: 1 counting A in the ATG of the START CODON as nucleotide position 1. The mutant cDNA is depicted in SEQ ID NO: 14.

[0060] "Mutant allele" refers herein to an allele comprising one or more mutations in the coding sequence (mRNA, cDNA or genomic sequence) compared to the wild type allele. Such mutation(s) (e.g. insertion, inversion, deletion and/or replacement of one or more nucleotide(s)) may lead to the encoded protein having reduced *in vitro* and/or *in vivo* functionality (reduced function) or no *in vitro* and/or *in vivo* functionality (loss-of-function), e.g. due to the protein e.g. being truncated or having an amino acid sequence wherein one or more amino acids are deleted, inserted or replaced. Such changes may lead to the protein having a different 3D conformation, being targeted to a different sub-cellular compartment, having a modified catalytic domain, having a modified binding activity to nucleic acids or proteins, etc.

[0061] "Wild type plant" and "wild type fruits" or "normal ripening" plants/fruits refers herein to a tomato plant comprising two copies of a wild type (WT) *Acs4* allele *(Acs4*/*Acs4)* encoding a fully functional Acs4 protein (e.g. in contrast to "mutant plants", comprising a mutant *acs4* allele). Such plants are for example suitable controls in phenotypic assays. Preferably wild type and/or mutant plants are "cultivated tomato plants". For example the cultivar Moneymaker is a wild type plant, as is cultivar Ailsa Craig, cultivar Tapa and many others.

[0062] "Tomato plants" or "cultivated tomato plants" are plants of the *Solanum lycopersicum,* i.e. varieties, breeding lines or cultivars of the species *Solanum lycopersicum,* cultivated by humans and having good agronomic characteristics; preferably such plants are not "wild plants", i.e. plants which generally have much poorer yields and poorer agronomic characteristics than cultivated plants and e.g. grow naturally in wild populations. "Wild plants" include for example ecotypes, PI (Plant Introduction) lines, landraces or wild accessions or wild relatives of a species. The so-called heirloom varieties or cultivars, i.e. open pollinated varieties or cultivars commonly grown during earlier periods in human history and often adapted to specific geographic regions, are in one aspect of the invention encompassed herein as cultivated tomato plants.

[0063] Wild relatives of tomato include *S. arcanum, S. chmielewskii, S. neorickii* (= *L. parviflorum), S. cheesmaniae, S. galapagense, S. pimpinellifolium, S. chilense, S. corneliomulleri, S. habrochaites* (= *L. hirsutum), S. huaylasense, S. sisymbriifolium, S. peruvianum, S. hirsutum* or *S. pennellii.*

[0064] "Average" refers herein to the arithmetic mean.

BRIEF DESCRIPTION OF THE SEQUENCE LISTING

[0065] SEQ ID NO: 1 shows the *Solanum lycopersicum* wild type, fully functional, ACS4 protein sequence as derived from the mRNA based on Genbank Accession number AAA34131.1 (encoded by the cDNA of GenBank Accession

number M63490.1).

[0066] SEQ ID NO: 2 shows the *Solanum lycopersicum* mutant 2477 acs4 protein sequence.

[0067] SEQ ID NO: 3 shows the *Solanum lycopersicum* mutant 4043 acs4 protein sequence.

[0068] SEQ ID NO: 4 shows the *Solanum lycopersicum* mutant 4222 acs4 protein sequence.

[0069] SEQ ID NO: 5 shows the *Solanum lycopersicum* mutant 4303 acs4 protein sequence.

[0070] SEQ ID NO: 6 shows the *Solanum lycopersicum* mutant 4691 acs4 protein sequence.

[0071] SEQ ID NO: 7 shows the *Solanum lycopersicum* mutant 5251 acs4 protein sequence.

[0072] SEQ ID NO: 8 shows the *Solanum lycopersicum* wild type *Acs4* cDNA based on Genbank Accession number M63490.1.

[0073] SEQ ID NO: 9 shows the *Solanum lycopersicum* mutant 2477 *acs4* cDNA.

[0074] SEQ ID NO: 10 shows the *Solanum lycopersicum* mutant 4043 *acs4* cDNA.

[0075] SEQ ID NO: 11 shows the *Solanum lycopersicum* mutant 4222 *acs4* cDNA.

[0076] SEQ ID NO: 12 shows the *Solanum lycopersicum* mutant 4303 *acs4* cDNA.

[0077] SEQ ID NO: 13 shows the *Solanum lycopersicum* mutant 4691 *acs4* cDNA.

[0078] SEQ ID NO: 14 shows the *Solanum lycopersicum* mutant 5251 *acs4* cDNA.

[0079] SEQ ID NO: 15 shows the *Solanum lycopersicum* wild type *Acs4* genomic DNA.

BRIEF DESCRIPTION OF THE FIGURES

[0080]

Figure 1: In this graph an alignment of the amino acid sequence of 5 of the sequences given in Figure 1 of Capitani et al. (Journal of Molecular Biology, 1999, vol 194, pp 745-756) (*Cucumis melo, Pelargonium hortorum, Brassica oleracea Phaseolus aureus,* and *Solanum tuberosum)* with the wild type *Solanum lycopersicum* ACS4 amino acid sequence as given in SEQ ID NO 1 is shown..

Figure 2: Ethylene-release measured in nl / (h · g), also written as $nl \cdot h^{-1} \cdot g^{-1}$, from tomato fruits at Pink stage and Red stage. Tapa is a commercial wild type cultivar *(Acs4*/*Acs4).*

Figure 3: In this graph the percentage of fruits in red stage is shown, determined at various days after the wild type control fruits started entering breaker stage [at day 1, the first fruit of Wild type was in breaker stage]. All fruits of mutant plants of the invention require more days to ripen compared to wild type (wt), 'Ho' means fruits of a mutant plant (indicated by the preceding number) being homozygous for a specific *acs4* mutation (*acs4*/*acs4*); He means fruits of a mutant (indicated by the preceding number) being heterozygous for a specific *acs4* mutation (*Acs4*/*acs4*).

Figure 4: Alignment of SEQ ID NO: 1 - 7. The Acs4 domains are also depicted (light gray), as are the mutations (in bold and underlined).

DETAILED DESCRIPTION OF THE INVENTION

[0081] The present invention discloses a cultivated plant of the species *Solanum lycopersicum* comprising an *acs4* allele having one or more mutations, said mutations resulting in production of a mutant acs4 protein having loss-of-function and / or reduced function compared to wild type Acs4 protein of SEQ ID NO: 1 wherein said mutation or mutations result in reduced ethylene production and/or delayed fruit ripening and/or a longer shelf life compared to *Solanum lycopersicum* being homozygous for the wild type ACS4 allele.

[0082] The Acs4 protein sequence contains 3 domains: a "large domain" referring to amino acid residues 65 to 327 as indicated in Figure 4 of this application and two small domains referring to amino acid residues 33 to 62 and 339 to 438, respectively as indicated in Figure 4 of this application. The Acs4 catalytic centre is believed to be in the "large domain".

[0083] In one aspect the invention relates to a cultivated plant of the species *Solanum lycopersicum,* and/or parts thereof (e.g. fruits), comprising an *acs4* allele having one or more mutations, said mutations resulting in production of a mutant acs4 protein having loss-of-function or reduced function compared to wild type Acs4 protein wherein said mutation or mutations result in reduced ethylene production and /or delayed fruit ripening and/or a longer shelf life compared to *Solanum lycopersicum* plants which are homozygous for the wild type fully functional *Acs4* allele (*Acs4*/*Acs4*) (encoding a functional Acs4 protein of SEQ ID NO: 1 or a functional variant).

[0084] A *S. lycopersicum* plant encoding the protein of SEQ ID NO: 1 is for example cultivar UC82B, or others.

[0085] In one aspect, a functional variant of SEQ ID NO: 1 is the Acs4 allele encoding the protein of GenBank accession CAH56694, CAH56504, or CAH56693. A S. *lycopersicum* plant encoding a functional variant of SEQ ID NO: 1 is for

example cultivar San Marzano Vesuvio, San Marzano Nano or Tondino.

[0086] In one aspect the invention relates to a cultivated plant of the species *Solanum lycopersicum,* and/or parts thereof (e.g. fruits), comprising an *acs4* allele having one or more mutations, said mutations resulting in production of a mutant acs4 protein having loss-of-function acs4 protein or reduced function compared to wild type Acs4 protein wherein said mutation or mutations result in reduced ethylene production and /or delayed fruit ripening and/or a longer shelf life compared to *Solanum lycopersicum* plants which are homozygous for the wild type fully functional *Acs4* allele *(Acs4*/*Acs4)* (encoding a functional Acs4 protein of SEQ ID NO: 1 or a functional variant), wherein the tomato plant does not comprise the Acs4 allele encoding the protein of GenBank accession CAH56694, CAH56504, or CAH56693. In another aspect, the mutation or mutations in the plant of the invention result in reduced ethylene production compared to *Solanum lycopersicum* being homozygous for the wild type *Acs4* allele.

[0087] In another aspect, the mutation or mutations in the plant of the invention result in delayed fruit ripening and/or a longer shelf life compared to *Solanum lycopersicum* being homozygous for the wild type *Acs4* allele.

[0088] In yet another aspect, the invention relates to a cultivated plant of the species *Solanum lycopersicum* comprising an *acs4* allele having one or more mutations resulting in a loss-of-function acs4 protein or reduced-function acs4 protein, wherein said mutation(s) are occurring in the "large domain", i.e. in the encoding part of amino acid region 65 to 327 of the wild type, functional Acs4 protein encoding, *Acs4* allele, and said mutations resulting in production of a mutant acs4 protein having loss-of-function acs4 protein or reduced function compared to wild type Acs4 protein wherein said mutation or mutations result in reduced ethylene production and/or delayed fruit ripening and/or a longer shelf life compared to *Solanum lycopersicum* being homozygous for the wild type *Acs4* allele. In a preferred aspect, the one or more mutations are one or more amino acid substitutions, deletions and/or insertions in the region of amino acid 241 to 251 of SEQ ID NO: 1 and/or in the region of amino acids 304 to 327 ; in another aspect the one or more mutations result in part or all of the large domain downstream of amino acid 200, 201, or 203 being absent or the mutations result in a truncated acs4 protein which lacks at least the second small domain and/or part of the large domain, e.g. a stop codon being present anywhere after nucleotide 600 of SEQ ID NO: 8.

[0089] In yet a further aspect, the invention relates to a *Solanum lycopersicum* plant comprising an *acs4* allele encoding a loss-of- function acs4 protein or reduced-function acs4 protein, which protein comprises a functional "large-domain", i.e. the mutation leading to the reduced ethylene production and/or the delayed ripening and/or longer shelf life, lies outside the "large-domain". Thus, in one embodiment the mutant *acs4* allele comprises one or more mutations in one or both of the small domains from amino acid 33 to 62 and/or 339 to 438 of SEQ ID NO:1, or of a variant of SEQ ID NO:1 which comprises a functional "large-domain", and further comprises (a nucleotide sequence encoding) at least one amino acid insertion, deletion or replacement in amino acids 33 to 62 and/or 339 to 438 of SEQ ID NO: 1, said at least one insertion, deletion or replacement leading to a reduced ethylene production and/or a delay in ripening and/or longer shelf life of the fruit of the tomato plant.

[0090] In one embodiment the mutation(s) causing the loss-of-function acs4 protein or reduced-function of the acs4 protein is/are in the "large-domain" of the wild type Acs4 protein i.e. which protein comprises functional "small-domains", thus in one embodiment one or more amino acids are inserted, deleted or replaced in amino acids 65 to 327 of SEQ ID NO: 1 or a variant of SEQ ID NO: 1. In another embodiment the mutation(s) causing the loss-of-function acs4 protein or reduced-function of the acs4 protein is/are in the C-terminus of the wild type Acs4 protein, thus in one embodiment one or more amino acids are inserted, deleted or replaced in amino acids 444 to 476 of SEQ ID NO: 1 (or a variant of SEQ ID NO: 1).

[0091] Thus, in one embodiment of the invention, the tomato plants according to the invention comprise an endogenous (non-transgenic) mutant *acs4* allele, which encodes a loss-of-function acs4 protein or reduced-function mutant acs4 protein whereby the fruits of the plant do ripen to the red stage (preferably slower than plants homozygous for the wild type allele, encoding a fully functional Acs4 protein). In another embodiment of the invention, the tomato plants according to the invention comprise a human-induced non-transgenic mutant *acs4* allele, which encodes a reduced-function mutant acs4 protein and/or a loss-of-function acs4 protein. In still another embodiment such mutant *acs4* allele is derived from and/or generated in a cultivated tomato (e.g. a breeding line, variety or heirloom variety) or a wild relative of tomato. Such a human-induced mutation may for example be induced using targeted mutagenesis as described in EP1963505. Mutant acs4 alleles generated in wild relatives of tomato are then easily transferred into cultivated tomato by breeding.

[0092] In still another aspect, the invention relates to a plant according to the invention having an endogenous *acs4* allele encoding a loss-of-function acs4 protein or reduced-function acs4 protein having substantial sequence identity to SEQ. ID NO: 1, or to a variant of SEQ ID NO: 1, wherein said protein comprising one or more amino acid replacements, deletions and/or insertions.

[0093] In yet another aspect, the invention relates to a plant of the invention comprising reduced ethylene production and/or delayed ripening and/or longer shelf-life than wild type *(Acs4*/*Acs4)* plants, due to said plants comprising an endogenous *acs4* allele encoding a loss-of-function acs4 protein or reduced-function acs4 protein having substantial sequence identity to SEQ. ID NO: 2 or to SEQ. ID NO: 3, or to SEQ. ID NO: 4, or to SEQ. ID NO: 5 or to SEQ. ID NO: 6, or to SEQ. ID NO: 7. In a specific aspect, the invention relates to cultivated tomato plants comprising a *acs4* allele as

found in seed deposited under accession number NCIMB 42034, NCIMB 42037, NCIMB 42038, NCIMB 42039, or NCIMB 42041 in one or two copies, i.e. in homozygous or heterozygous form. In heterozygous form, the other allele may be a wild type *Acs4* allele or another mutant *acs4* allele, such as from any one of the other mutants provided herein, or any other mutant *acs4* allele encoding for a loss-of-function acs4 protein or reduced-function acs4 protein as described herein. In heterozygous form, the other allele may thus be a reduced function *acs4* allele.

[0094] In still another aspect, the invention relates to a an endogenous *acs4* allele or to a loss-of-function acs4 protein or a reduced-function acs4 protein encoded by it having substantial sequence identity to SEQ. ID NO: 2 or to SEQ. ID NO: 3, or to SEQ. ID NO: 4, or to SEQ. ID NO: 5 or to SEQ. ID NO: 6, or to SEQ. ID NO: 7 as found in (and as derivable from) seed deposited under accession number NCIMB 42034, NCIMB 42037, NCIMB 42038, NCIMB 42039, orNCIMB 42041.

[0095] In yet another aspect, the invention relates to a tomato plant of the invention comprising an endogenous *acs4* allele encoding a loss-of-function acs4 protein or reduced-function acs4 protein having 100% sequence identity to SEQ. ID NO: 2, or to SEQ. ID NO: 3, or to SEQ. ID NO: 4, or to SEQ. ID NO: 5, or to SEQ. ID NO: 6, or to SEQ. ID NO: 7.

[0096] In yet a further aspect, the invention relates to a plant of the invention comprising an endogenous *acs4* allele encoding a loss-of-function acs4 protein or reduced-function acs4 protein having at least one amino acid deletion, insertion or replacement in the "large-domain". Preferably the acs4 protein comprises functional small domains, such as the small domains of SEQ ID NO: 1 (acid residues 33 to 62 and/or 339 to 438) or the small domains of a (functional) variant of SEQ ID NO: 1. In one embodiment it also comprises the C-terminal of SEQ ID NO: 1 (amino acids 444 to 476) or the C-terminal of a (functional) variant of SEQ ID NO: 1.

[0097] In one aspect, the acs4 protein is not longer than 203 amino acids preferably the first 203 amino acids. Thus, in one embodiment the tomato plant encodes a truncated acs4 protein, comprising amino acids 1-450, 1-400, 1-350, 1-300, 1-250, or 1-203 of SEQ ID NO: 1 or a variant thereof.

[0098] The invention further relates to tomato seeds, plants and plant parts comprising an endogenous *acs4* gene encoding a cDNA (mRNA) having substantial sequence identity to SEQ. ID NO: 8 and having at least one non-transgenic mutation within said endogenous *acs4* gene, wherein said at least one non-transgenic mutation results in the production of a mutant acs4 protein having loss-of-function acs4 protein or reduced activity compared to wild type Acs4 protein. Preferably, said mutation results in reduced ethylene production and/or slower fruit ripening and/or a longer shelf life compared to *Solanum lycopersicum* being homozygous for the functional wild type *Acs4* allele, encoding the protein of SEQ ID NO: 1 or a functional variant thereof. The mutation described anywhere herein may be human-induced or it may be a natural mutation. The plant is preferably a cultivated tomato plant. In one embodiment the mutation results in either a stop-codon or in an amino acid substitution. In one embodiment the amino acid selected from the group consisting of Ala248, Val250, Ser279, Thr316 and Leu321 of the wild type Acs4 protein is substituted for a different amino acid, e.g. Ala248Val, Val250Glu, Ser279Asn, Thr316Ile and Leu321Phe. In another embodiment, said mutation is selected from the group consisting of G836A, C743T, A610T, G963T, T749A, and C947T of SEQ ID NO: 8.

[0099] In another aspect the invention relates to tomato seeds, plants and plant parts comprising an endogenous mutant *acs4* gene wherein said non-transgenic mutation creates an amino acid change in the acs4 protein encoded by and produced by transcription and translation of the *acs4* gene, wherein said amino acid change is selected from the group consisting of S279N, A248V, L321F, V250E, T316I, and the complete deletion of amino acids 204 to 476 of SEQ ID NO: 1.

[0100] In yet another aspect the invention relates to acs4 protein having substantial sequence identity to SEQ ID NO: 2. In still another aspect the invention relates to acs4 protein having substantial sequence identity to SEQ ID NO: 3. In a further aspect the invention relates to acs4 protein having substantial sequence identity to SEQ ID NO: 4. In yet another aspect the invention relates to acs4 protein having substantial sequence identity to SEQ ID NO: 5. In still another aspect the invention relates to acs4 protein having substantial sequence identity to SEQ ID NO: 6. In a further aspect the invention relates to acs4 protein having substantial sequence identity to SEQ ID NO: 7. The invention also relates to tomato seeds, plants and plant parts comprising a nucleotide sequence encoding these proteins.

[0101] In still another aspect, the invention relates to tomato fruit, seeds, pollen, plant parts, and/or progeny of a plant of the invention. Preferably, the invention relates to fruit or seeds of the plant of the invention. More preferably, the invention relates to tomato fruit having delayed ripening and/or an increased post-harvest shelf life caused by a non-transgenic mutation in at least one *acs4* allele, as described elsewhere herein.

[0102] In one aspect the tomato plants according to the invention have a delay of breaker stage, meaning that the mutants according to the invention require significantly more days e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 or more days than wild type *Acs4*/*Acs4* controls for the first fruits and/or for all fruits to have entered breaker stage.

[0103] In another aspect fruits of the tomato plants of the invention require more days to go from breaker stage to red stage , e.g. fruits of the plants of the invention require 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 or more days than wild type *Acs4*/*Acs4* controls to go from breaker stage to red stage.

[0104] In another aspect the invention relates to a fruit of a plant of the invention having a the shelf life that is at least 2 days longer than the shelf life of a tomato fruit being homozygous for the wild type Acs4 allele. In still another aspect

the invention relates to a fruit according to a plant of the invention having a reduced ethylene production that is at least 15% reduced, or at least 20 % reduced compared to *Solanum lycopersicum* being homozygous for the wild type Acs4 allele.

[0105] In a particular aspect the tomato plants according to the invention have a shelf life that is significantly longer than the shelf life of wild type plants, for example the number of days from the first fruit being in breaker stage (or turning stage, pink stage, red stage or from harvest) up to the first fruit starting to become 'bad' and unsuitable for sale or consumption is significantly longer, e.g. at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more, days longer than fruits of control plants (such as wild type *Acs4*/*Acs4* plants), when plants are grown under the same conditions and fruits are treated the same way and kept under the same conditions.

[0106] A delayed ripening and/or extended shelf-life can have the advantage that more time is available for transport of picked fruits e.g. to retailers and supermarkets and/or that the consumer can keep the fruits longer. Tomatoes can be harvested at mature green stage or at breaker stage, or thereafter. When harvested before breaker stage, ethylene exposure is needed, while harvest around breaker stage or thereafter does not require ethylene exposure, as the fruits produce ethylene themselves. As seen in Figure 2, delayed-ripening mutants according to the invention produce less ethylene at pink stage and red stage than wild type fruits, but sufficient ethylene to ripen to the red stage. In one aspect of the invention tomato plants are provided comprising a mutant *acs4* allele encoding a loss-of-function acs4 protein or reduced function acs4 protein, wherein the fruits of said plants produce significantly less ethylene than wild type *(Acs4*/*Acs4)* plants. "Significantly less ethylene" refers to the fruit producing equal to or less than 75%, equal to or less than 70%, equal to or less than 65%, equal to or less than 60%, equal to or less than 55%, equal to or less than 50%, equal to or less than 45%, equal to or less than 40%, equal to or less than 35%, equal to or less than 30%, equal to or less than 25% equal to or less than 20%, or equal to or less than 15% of the ethylene produced by homozygeous *Acs4*/*Acs4* fruits at the pink or red stage. Thus, the ethylene produced at the pink stage is in one aspect below about 3.5 nl/(h · g), such as equal to or below about 3 nl/(h · g) or equal to or below about 2.5 nl/(h · g) or equal to or below about 2.0 nl/(h · g) or equal to or below about 1.5 nl/(h · g) or equal to or below about 1.0 nl/(h · g)or equal to or below about 0.5 nl/(h · g). The ethylene produced at the red stage is in one aspect below about 6 nl/(h · g), such as equal to or below about 5.5 nl/(h · g) or equal to or below about 5.0 nl/(h · g), or equal to or below 4.5 nl/(h · g), or equal to or below about 3.5 nl/(h · g), or equal to or below about 3 nl/(h · g) or equal to or below about 2.5 nl/(h · g) or equal to or below about 2.0 nl/(h · g) or equal to or below about 1.5 nl/(h · g) or equal to or below about 1.0 nl/(h · g)or equal to or below about 0.5 nl/(h · g).

[0107] In another aspect, the invention relates to tomato fruit of a plant of the invention having a longer ripening period and/or an increased post-harvest shelf life caused by a non-transgenic mutation in at least one *acs4* allele wherein the longer ripening period and/or the longer post-harvest shelf life is at least 110% of the ripening period and/or of the post-harvest shelf life of a tomato fruit being homozygous for the wild type *Acs4* allele. Preferably, the ripening period and/or post-harvest shelf life is at least 115%, more preferably at least 120%, even more preferably at least 125% of the ripening period and/or post-harvest shelf life of a tomato fruit being homozygous for the wild type *Acs4* allele. In another aspect, the ripening period and/or post-harvest shelf life is at least 135%, more preferably at least 150%, even more preferably at least 165% of the ripening period and/or post-harvest shelf life of a tomato fruit being homozygous for the wild type *Acs4* allele. In yet another aspect, the ripening period and/or post-harvest shelf life is at least 180%, more preferably at least 200% even more preferably at least 250 % of the ripening period and/or post-harvest shelf life of a tomato fruit being homozygous for the wild type *Acs4* allele.

[0108] In still another aspect of the invention tomato plants are provided that have the same or similar delayed ripening and/or increased shelf life as tomato plants of the invention, of which representative seeds were deposited by Nunhems B.V. and accepted for deposit on 21 August 2012 at the NCIMB Ltd. (Ferguson Building, Craibstone Estate, Bucksburn Aberdeen, Scotland AB21 9YA, UK) according to the Budapest Treaty, under the Expert Solution (EPC 2000, Rule 32(1)). Seeds were given the following deposit numbers: NCIMB 42034 (mutant 2477), NCIMB 42037 (mutant 4043), NCIMB 42038 (mutant 4222), NCIMB 42039 (mutant 4691), NCIMB 42041 (mutant 5251).

[0109] According to a further aspect the invention provides a cell culture or tissue culture of the tomato plant of the invention. The cell culture or tissue culture comprises regenerable cells. Such cells can be derived from leaves, pollen, embryos, cotyledon, hypocotyls, meristematic cells, roots, root tips, anthers, flowers, seeds and stems.

[0110] Seeds from which plants according to the invention can be grown are also provided, as well as packages containing such seeds. Also a vegetative propagation of plants according to the invention are an aspect encompassed herein. Likewise harvested fruits and fruit parts, either for fresh consumption or for processing or in processed form are encompassed. Fruits may be graded, sized and/or packaged. Fruits may be sliced or diced or further processed.

[0111] In another aspect the invention relates to one or more cells of a plant of the invention.

[0112] Also described are food and/ or food products incorporating the fruit or part of a fruit of a tomato plant of the invention. As used herein, food refers to nutrients consumed by human or animal species. Examples are sandwiches, salads, sauces, ketchup and the like.

[0113] Also described is a method of producing a tomato plant of the invention comprising the steps of:

a. obtaining plant material from a tomato plant;
b. treating said plant material with a mutagen to create mutagenized plant material;
c. analyzing said mutagenized plant material to identify a plant having at least one mutation in at least one *acs4* allele having substantial sequence identity to SEQ ID NO: 1 or variants thereof.

[0114] The method may further comprise analyzing the ripening period and/or shelf life of tomato fruits of the selected plant or progeny of the plant and selecting a plant of which the fruit have delayed ripening and/or extended shelf-life. The mutation may be selected from a mutation in the large domain of the acs4 protein and/or in the second small domain of the acs4 protein (amino acids 339-438). In one aspect the mutation is selected from a mutation resulting in an amino acid substitution selected from the group consisting of S279N, A248V, L321F, V250E, T316I, or from a stop-codon mutation causing the deletion of amino acids 204 to 476 of SEQ ID NO: 1 or of a part thereof . In a further aspect, the mutation is selected from a mutation cuasing a change in the cDNA selected from the group G836A, C743T, A610T, G963T, T749A, and C947T of SEQ ID NO: 8.. In this method, the plant material of step a) is preferably selected from the group consisting of seeds, pollen, plant cells, or plant tissue of a tomato plant line or cultivar. Plant seeds being more preferred. The mutagen used in this method may be ethyl methanesulfonate. In step b) and step c) the mutagenized plant material is preferably a mutant population, such as a tomato TILLING population.

[0115] Thus, a method for producing a tomato plant comprising delayed fruit ripening and/or longer fruit shelf-life is described comprising the steps of:

a) providing a tomato TILLING population,

b) screening said TILLING population for mutants in the *acs4* gene, especially in the large-domain encoding nucleotide sequence, and

c) selecting from the mutant plants of b) those plants (or progeny of those plants) of which the fruits have a reduced ethylene production and/or a delayed ripening and/or longer shelf life than wild type *(Acs4*/*Acs4)* fruits.

[0116] Mutant plants (M1) are preferably selfed one or more times to generate for example M2 populations or preferably M3 or M4 populations for phenotyping. In M2 populations the mutant allele is present in a ratio of 1 (homozygous for mutant allele) : 2 (heterozygous for mutant allele): 1 (homozygous for wild type allele).

[0117] Also described is a method for producing a hybrid *Solanum lycopersicum* plant, said method comprising:

(a) obtaining a first *Solanum lycopersicum* plant of the current invention and

(b) crossing said first *Solanum lycopersicum* plant with a second *Solanum lycopersicum* plant;

wherein said hybrid *Solanum lycopersicum* plant comprises an *acs4* allele having one or more mutations wherein said mutations result in production of a mutant acs4 protein having loss-of-function acs4 protein or reduced activity compared to wild type Acs4 protein.

[0118] Plants and plant parts (e.g. fruits, cells, etc.) of the invention can homozygous or heterozygous for the mutant *acs4* allele.

[0119] Preferably the plants according to the invention, which comprise one or more mutant *acs4* alleles (or variants), and which produce a mutant acs4 protein having loss-of-function acs4 protein or reduced activity compared to wild type Acs4 protein, do not produce fewer fruits than the wild type plants. Thus, fruit number per plant is preferably not reduced.

[0120] Other putative ACS4 genes/proteins can be identified *in silico,* e.g. by identifying nucleic acid or protein sequences in existing nucleic acid or protein database (e.g. GENBANK, SWISSPROT, TrEMBL) and using standard sequence analysis software, such as sequence similarity search tools (BLASTN, BLASTP, BLASTX, TBLAST, FASTA, etc.).

[0121] In one embodiment loss-of-function acs4 protein or reduced-function mutant acs4 proteins (including variants or orthologs, such as acs4 proteins of wild tomato relatives) are provided and plants and plant parts comprising one or more *acs4* alleles in their genome, which encode loss-of-function acs4 protein or reduced-function mutants, whereby the reduced-function confers reduced ethylene production and /or slower fruit ripening and/or a longer shelf life compared to *Solanum lycopersicum* being homozygous for the wild type *Acs4* allele.

[0122] Any type of mutation may lead to a reduction in function of the encoded Acs4 protein, e.g. insertion, deletion and/or replacement of one or more nucleotides in the genomic DNA which comprises the cDNA (SEQ ID NO: 8, or variants thereof). In a preferred embodiment is provided an *acs4* nucleic acid sequence capable of reduced ethylene production and/or conferring slower fruit ripening and/or a longer shelf life compared to *Solanum lycopersicum* being homozygous for the wild type *Acs4* allele, whereby the nucleic acid sequence encodes a loss-of-function acs4 protein

or reduced-function Acs4 protein due to one or more mutations in the large domain.

[0123] The *in vivo* loss-of-function acs4 protein or reduced-function of such proteins can be tested as described herein, by determining the effect this mutant allele has on ethylene production and/or ripening period and/or shelf life period. Plants comprising a nucleic acid sequence encoding such mutant loss-of-function acs4 protein or reduced-function proteins and having a reduced ethylene production and/or slower fruit ripening and/or a longer shelf life compared to *Solanum lycopersicum* being homozygous for the wild type *Acs4* allele can for example be generated using e.g. mutagenesis and identified by TILLING or identified using EcoTILLING, as known in the art. Also transgenic methods can be used to test *in vivo* functionality of a mutant *acs4* allele encoding a mutant acs4 protein. A mutant allele can be operably linked to a plant promoter and the chimeric gene can be introduced into a tomato plant by transformation. Regenerated plants (or progeny, e.g. obtained by selfing), can be tested for ethylene production and/or fruit ripening period and/or shelf life. For example a tomato plant comprising a non-functional *acs4* allele can be transformed to test the functionality of the transgenic *acs4* allele.

[0124] TILLING (Targeting Induced Local Lesions IN Genomes) is a general reverse genetic technique that uses traditional chemical mutagenesis methods to create libraries of mutagenized individuals that are later subjected to high throughput screens for the discovery of mutations. TILLING combines chemical mutagenesis with mutation screens of pooled PCR products, resulting in the isolation of missense and non-sense mutant alleles of the targeted genes. Thus, TILLING uses traditional chemical mutagenesis (e.g. EMS or MNU mutagenesis) or other mutagenesis methods (e.g. radiation such as UV) followed by high-throughput screening for mutations in specific target genes, such as *Acs4* according to the invention. S1 nucleases, such as CEL1 or ENDO1, are used to cleave heteroduplexes of mutant and wildtype target DNA and detection of cleavage products using e.g. electrophoresis such as a LI-COR gel analyzer system, see e.g. Henikoff et al. Plant Physiology 2004, 135: 630-636. TILLING has been applied in many plant species, such as tomato. (see http://tilling.ucdavis.edu/index.php/Tomato_Tilling), rice (Till et al. 2007, BMC Plant Biol 7: 19), Arabidopsis (Till et al. 2006, Methods Mol Biol 323: 127-35),-Brassica, maize (Till et al. 2004, BMC Plant Biol 4: 12), etc. Also EcoTILLING, whereby mutants in natural populations are detected, has been widely used, see Till et al. 2006 (Nat Protoc 1: 2465-77) and Comai et al. 2004 (Plant J 37: 778-86).

[0125] In one embodiment of the invention (cDNA or genomic) nucleic acid sequences encoding such mutant acs4 proteins comprise one or more non-sense and/or missense mutations, e.g. transitions (replacement of purine with another purine (A ↔ G) or pyrimidine with another pyrimidine (C ↔ T)) or transversions (replacement of purine with pyrimidine, or *vice versa* (C/T ↔ A/G). In one embodiment the non-sense and/or missense mutation(s) is/are in the nucleotide sequence encoding any of the Acs4 exons, more preferably in the ACS4 large domain or an essentially similar domain of a variant Acs4 protein, i.e. in a domain comprising at least 80%, 90%, 95%, 98%, 99% amino acid identity to amino acids 65-327 of SEQ ID NO: 1 or variants thereof.

[0126] In one embodiment an *acs4* nucleotide sequence comprising one or more non-sense and/or missense mutations in one of the exon- encoding sequence are provided, as well as a plant comprising such a mutant allele resulting in reduced ethylene production and/or delayed fruit ripening and/or a longer shelf life compared to *Solanum lycopersicum* being homozygous for the wild type *Acs4* allele.

[0127] In a specific embodiment of the invention tomato plants and plant parts (fruits, seeds, etc.) comprising a mutant loss-of-function or reduced-function *acs4* allele are provided.

[0128] In one embodiment, the loss-of-function acs4 protein or reduced-function acs4 protein is a truncated protein, i.e. a protein fragment of any one of the Acs4 proteins defined further above (including variants thereof). In general EMS (Ethyl methanesulfonate) induces substitutions of guanine/cytosine to adenin/thymine. In case of a glutamine (Gln or Q, encoded by the nucleotides CAA or CAG) or arginine (Arg or R, encoded by the nucleotides CGA) codon, a substitution of the cytosine for thymine can lead to the introduction of a stop codon in the reading frame (for example CAA/CAG/CGA to TAA/TAG/TGA) resulting in a truncated protein.

[0129] Also provided are nucleic acid sequences (genomic DNA, cDNA, RNA) encoding loss-of-function acs4 protein or reduced-function acs4 proteins, such as for example acs4 depicted in SEQ ID NO: 2, 3, 4, 5, 6, or 7; or variants thereof as defined above (including any chimeric or hybrid proteins or mutated proteins or truncated proteins). Due to the degeneracy of the genetic code various nucleic acid sequences may encode the same amino acid sequence. The nucleic acid sequences provided include naturally occurring, artificial or synthetic nucleic acid sequences. A nucleic acid sequence encoding Acs4 is provided for in SEQ ID NO: 8 (wild type cDNA), Genbank Accession Number M63490.1.

[0130] It is understood that when sequences are depicted as DNA sequences while RNA is referred to, the actual base sequence of the RNA molecule is identical with the difference that thymine (T) is replace by uracil (U). When referring herein to nucleotide sequences (e.g DNA or RNA) italics are used, e.g. *acs4* allele, while when referring to proteins, no italics are used, e.g. acs4 protein. Mutants are in small letters (e.g *acs4* allele or acs4 protein), while wild type / functional forms start with a capital letter *(Acs4* allele or Acs4 protein).

[0131] Also provided are nucleic acid sequences (genomic DNA, cDNA, RNA) encoding mutant acs4 proteins, i.e. loss-of-function acs4 protein or reduced function acs4 proteins, as described above, and plants and plant parts comprising such mutant sequences. For example, *acs4* nucleic acid sequences comprising one or more non-sense and/or missense

mutations in the wild type *Acs4* coding sequence, rendering the encoded protein having a loss-of-function or reduced function *in vivo.* Also sequences with other mutations are provided, such as splice-site mutants, i.e. mutations in the genomic *acs4* sequence leading to aberrant splicing of the pre-mRNA, and/or frame-shift mutations, and/or insertions (e.g. transposon insertions) and/or deletions of one or more nucleic acids.

[0132] It is clear that many methods can be used to identify, synthesise or isolate variants or fragments of *acs4* nucleic acid sequences, such as nucleic acid hybridization, PCR technology, *in silico* analysis and nucleic acid synthesis, and the like. Variants of SEQ ID NO: 8, may either encode wild type, functional Acs4 proteins, or they may encode loss-of-function acs4 protein or reduced-function mutant alleles of any of these, as for example generated e.g. by mutagenesis and/or identified by methods such as TILLING or EcoTILLING, or other methods.

[0133] A plant of the invention can be used in a conventional plant breeding scheme to produce more plants with the same characteristics or to introduce the mutated *acs4* allele into other plant lines or varieties of the same or related plant species.

[0134] Also transgenic plants can be made using the mutant *acs4* nucleotide sequences of the invention using known plant transformation and regeneration techniques in the art. An "elite event" can be selected, which is a transformation event having the chimeric gene (comprising a promoter operably linked to a nucleotide sequence encoding a loss-of-function acs4 protein or reduced-function acs4 protein) inserted in a particular location in the genome, which results in good expression of the desired phenotype.

[0135] The plants of the invention as described above are homozygous for the mutant *acs4* allele, or heterozygous. To generate plants comprising the mutant allele in homozygous form, selfing can be used. The mutant *acs4* alleles according to the invention can be transferred to any other tomato plant by traditional breeding techniques, such as crossing, selfing, backcrossing, etc. Thus any type of tomato having delayed ripening and/or longer shelf life due to the presence of at least one mutant *acs4* allele according to the invention can be generated. Any *S. lycopersicum* may be generated and/or identified having at least one mutant *acs4* allele in its genome and producing a acs4 protein having loss-of-function acs4 protein or reduced activity compared to wild type Acs4 protein. The tomato plant may, thus, be any cultivated tomato, any commercial variety, any breeding line or other, it may be determinate or indeterminate, open pollinated or hybrid, producing fruits of any colour, shape and size. The mutant allele generated and/or identified in a particular tomato plant, or in a sexually compatible relative of tomato, may be easily transferred into any other tomato plant by breeding (crossing with a plant comprising the mutant allele and then selecting progeny comprising the mutant allele).

[0136] The presence or absence of a mutant *acs4* allele according to the invention in any tomato plant or plant part and/or the inheritance of the allele to progeny plants can be determined phenotypically and/or using molecular tools (e.g. detecting the presence or absence of the *acs4* nucleotide or acs4 protein using direct or indirect methods).

[0137] The mutant allele is in one embodiment generated or identified in a cultivated plant, but may also be generated and/or identified in a wild plant or non-cultivated plant and then transferred into an cultivated plant using e.g. crossing and selection (optionally using interspecific crosses with e.g. embryo rescue to transfer the mutant allele). Thus, a mutant *acs4* allele may be generated (human induced mutation using mutagenesis techniques to mutagenize the target *acs4* gene or variant thereof) and/or identified (spontaneous or natural allelic variation) in *Solanum lycopersicum* or in other *Solanum* species include for example wild relatives of tomato, such as S. *cheesmanii,* S. *chilense, S. habrochaites (L. hirsutum), S. chmielewskii, S. lycopersicum x S. peruvianum,* S. *glandulosum,* S. *hirsutum, S. minutum, S. parviflorum, S. pennellii,* S. *peruvianum,* S. *peruvianum* var. *humifusum* and *S. pimpinellifolium,* and then transferred into a cultivated *Solanum* plant, e.g. *Solanum lycopersicum* by traditional breeding techniques. The term "traditional breeding techniques" encompasses herein crossing, selfing, selection, double haploid production, embryo rescue, protoplast fusion, transfer via bridge species, etc. as known to the breeder, i.e. methods other than genetic modification by which alleles can be transferred.

[0138] The plant comprising the mutant *acs4* allele (e.g. tomato) may be crossed with another plant of the same species or of a closely related species, to generate a hybrid plant (hybrid seed) comprising the mutant *acs4* allele. Such a hybrid plant is also an embodiment of the invention.

[0139] In one embodiment F1 hybid tomato seeds (i.e. seeds from which F1 hybrid tomato plants can be grown) are provided, comprising at least one *acs4* allele according to the invention. F1 hybrid seeds are seeds harvested from a cross between two inbred tomato parent plants. Such an F1 hybrid may comprise one or two mutant *acs4* alleles according to the invention. Thus, in one embodiment a plant according to the invention is used as a parent plant to produce an F1 hybrid, the fruit of which have reduced ethylene production and/or delayed ripening and/or longer shelf-life than wild type *Acs4*/*Acs4* plants.

[0140] Also a method for transferring a mutant *acs4* allele to another plant is described, comprising providing a plant comprising a mutant *acs4* allele in its genome, whereby the mutant allele produce fruits that show reduced ethylene production and/or slower fruit ripening and/or a longer shelf life compared to *Solanum lycopersicum* being homozygous for the wild type *Acs4* allele (as described above), crossing said plant with another plant and obtaining the seeds of said cross. Optionally plants obtained from these seeds may be further selfed and/or crossed and progeny selected comprising

the mutant allele and producing fruits with delayed ripening and/or longer shelf-life and / or reduced ethylene production due to the presence of the mutant allele compared to plants comprising the wild type *Acs4* allele.

[0141] As mentioned, it is understood that other mutagenesis and/or selection methods may equally be used to generate mutant plants according to the invention. Seeds may for example be radiated or chemically treated to generate mutant populations. Also direct gene sequencing of *acs4* may be used to screen mutagenized plant populations for mutant alleles. For example KeyPoint screening is a sequence based method which can be used to identify plants comprising mutant *acs4* alleles (Rigola et al. PloS One, March 2009, Vol 4(3):e4761).

[0142] Thus, non-transgenic mutant tomato plants which produce lower levels of wild type Acs4 protein in fruits are provided, or which completely lack wild type Acs4 protein in fruits, and which produce loss-of- function acs4 protein or reduced-function acs4 protein in fruits due to one or more mutations in one or more endogenous *acs4* alleles, are provided. These mutants may be generated by mutagenesis methods, such as TILLING or variants thereof, or they may be identified by EcoTILLING or by any other method. *Acs4* alleles encoding loss-of- function acs4 protein or reduced-functional acs4 protein may be isolated and sequenced or may be transferred to other plants by traditional breeding methods.

[0143] Any part of the plant, or of the progeny thereof, is provided, including harvested fruit, harvested tissues or organs, seeds, pollen, flowers, ovaries, etc. comprising a mutant *acs4* allele according to the invention in the genome. Also plant cell cultures or plant tissue cultures comprising in their genome a mutant *acs4* allele are provided. Preferably, the plant cell cultures or plant tissue cultures can be regenerated into whole plants comprising a mutant *acs4* allele in its genome. Also double haploid plants (and seeds from which double haploid plants can be grown), generated by chromosome doubling of haploid cells comprising an *acs4* mutant allele, and hybrid plants (and seeds from which hybrid plants can be grown) comprising a mutant *acs4* allele in their genome are encompassed herein, whereby the double haploid plants and hybrid plants produce delayed ripening and/or longer shelf life fruits according to the invention.

[0144] Preferably, the mutant plants also have good other agronomic characteristics, i.e. they do not have reduced fruit numbers and/or reduced fruit quality compared to wild type plants. In a preferred embodiment the plant is a tomato plant and the fruit is a tomato fruit, such as a processing tomato, fresh market tomato of any shape or size or colour. Thus, also harvested products of plants or plant parts comprising one or two mutant *acs4* alleles are provided. This disclosure includes downstream processed products, such as tomato paste, ketchup, tomato juice, cut tomato fruit, canned fruit, dried fruit, peeled fruit, etc. The products can be identified by comprising the mutant allele in their genomic DNA.

Seed Deposits

[0145] A representative sample of seeds of five tomato TILLING mutants according to Example 1, were deposited by Nunhems B.V. and accepted for deposit on 21 August 2012 at the NCIMB Ltd. (Ferguson Building, Craibstone Estate, Bucksburn Aberdeen, Scotland AB21 9YA, UK) according to the Budapest Treaty, under the Expert Solution (EPC 2000, Rule 32(1)). Seeds were given the following deposit numbers: NCIMB 42034 (mutant 2477), NCIMB 42037 (mutant 4043), NCIMB 42038 (mutant 4222), NCIMB 42039 (mutant 4691), NCIMB 42041 (mutant 5251).

[0146] The Applicant requests that samples of the biological material and any material derived therefrom be only released to a designated Expert in accordance with Rule 32(1) EPC or related legislation of countries or treaties having similar rules and regulation, until the mention of the grant of the patent, or for 20 years from the date of filing if the application is refused, withdrawn or deemed to be withdrawn.

[0147] Access to the deposit will be available during the pendency of this application to persons determined by the Director of the U.S. Patent Office to be entitled thereto upon request. Subject to 37 C.F.R. § 1.808(b), all restrictions imposed by the depositor on the availability to the public of the deposited material will be irrevocably removed upon the granting of the patent. The deposit will be maintained for a period of 30 years, or 5 years after the most recent request, or for the enforceable life of the patent whichever is longer, and will be replaced if it ever becomes nonviable during that period. Applicant does not waive any rights granted under this patent on this application or under the Plant Variety Protection Act (7 USC 2321 et seq.).

SEQUENCE LISTING

[0148]

<110> Nunhems B.V.

<120> Solanum lycopersicum plants having non-transgenic alterations in the acs4 gene

<130> BCS 12-8012

<160> 15

<170> PatentIn version 3.5

<210> 1
<211> 476
<212> PRT
<213> Lycopersicon esculentum

<400> 1

```
Met Asp Leu Glu Thr Ser Glu Ile Ser Asn Tyr Lys Ser Ser Val Val
1               5                  10                  15

Leu Ser Lys Leu Ala Ser Asn Glu Gln His Gly Glu Asn Ser Pro Tyr
            20                  25                  30

Phe Asp Gly Trp Lys Ala Tyr Asp Asn Asp Pro Phe His Leu Val Asn
        35                  40                  45

Asn Leu Asn Gly Val Ile Gln Met Gly Leu Ala Glu Asn Gln Leu Ser
    50                  55                  60

Val Asp Leu Ile Glu Glu Trp Ile Lys Arg Asn Pro Lys Ala Ser Ile
65                  70                  75                  80

Cys Thr Asn Asp Gly Ile Glu Ser Phe Arg Arg Ile Ala Asn Phe Gln
                85                  90                  95

Asp Tyr His Gly Leu Pro Glu Phe Thr Asn Ala Ile Ala Lys Phe Met
            100                 105                 110

Glu Lys Thr Arg Gly Gly Lys Val Lys Phe Asp Ala Lys Arg Val Val
        115                 120                 125

Met Ala Gly Gly Ala Thr Gly Ala Asn Glu Thr Leu Ile Leu Cys Leu
    130                 135                 140

Ala Asp Pro Gly Asp Ala Phe Leu Val Pro Thr Pro Tyr Tyr Pro Gly
145                 150                 155                 160

Phe Asn Arg Asp Leu Arg Trp Arg Ser Gly Val Gln Leu Leu Pro Ile
                165                 170                 175
```

```
Ser Cys Lys Ser Cys Asn Asn Phe Lys Ile Thr Ile Glu Ala Ile Glu
            180                 185                 190

Glu Ala Tyr Glu Lys Gly Gln Gln Ala Asn Val Lys Ile Lys Gly Leu
        195                 200                 205

Ile Leu Thr Asn Pro Cys Asn Pro Leu Gly Thr Ile Leu Asp Arg Asp
    210                 215                 220

Thr Leu Lys Lys Ile Ser Thr Phe Thr Asn Glu His Asn Ile His Leu
225                 230                 235                 240

Val Cys Asp Glu Ile Tyr Ala Ala Thr Val Phe Asn Ser Pro Lys Phe
                245                 250                 255

Val Ser Ile Ala Glu Ile Ile Asn Glu Asp Asn Cys Ile Asn Lys Asp
            260                 265                 270

Leu Val His Ile Val Ser Ser Leu Ser Lys Asp Leu Gly Phe Pro Gly
        275                 280                 285

Phe Arg Val Gly Ile Val Tyr Ser Phe Asn Asp Asp Val Val Asn Cys
    290                 295                 300

Ala Arg Lys Met Ser Ser Phe Gly Leu Val Ser Thr Gln Thr Gln His
305                 310                 315                 320

Leu Leu Ala Phe Met Leu Ser Asp Asp Glu Phe Val Glu Glu Phe Leu
                325                 330                 335

Ile Glu Ser Ala Lys Arg Leu Arg Glu Arg Tyr Glu Lys Phe Thr Arg
            340                 345                 350

Gly Leu Glu Glu Ile Gly Ile Lys Cys Leu Glu Ser Asn Ala Gly Val
        355                 360                 365

Tyr Cys Trp Met Asp Leu Arg Ser Leu Leu Lys Glu Ala Thr Leu Asp
    370                 375                 380

Ala Glu Met Ser Leu Trp Lys Leu Ile Ile Asn Glu Val Lys Leu Asn
385                 390                 395                 400

Val Ser Pro Gly Ser Ser Phe Asn Cys Ser Glu Val Gly Trp Phe Arg
                405                 410                 415

Val Cys Phe Ala Asn Ile Asp Asp Gln Thr Met Glu Ile Ala Leu Ala
            420                 425                 430
```

```
Arg Ile Arg Met Phe Met Asp Ala Tyr Asn Asn Val Asn Lys Asn Gly
        435                 440                 445

Val Met Lys Asn Lys His Asn Gly Arg Gly Thr Thr Tyr Asp Leu Thr
    450                 455                 460

Pro Gln Met Gly Ser Thr Met Lys Met Leu Leu Ala
465                 470                 475
```

<210> 2
<211> 476
<212> PRT
<213> Lycopersicon esculentum

<400> 2

```
Met Asp Leu Glu Thr Ser Glu Ile Ser Asn Tyr Lys Ser Ser Val Val
1               5                   10                  15

Leu Ser Lys Leu Ala Ser Asn Glu Gln His Gly Glu Asn Ser Pro Tyr
            20                  25                  30

Phe Asp Gly Trp Lys Ala Tyr Asp Asn Asp Pro Phe His Leu Val Asn
        35                  40                  45

Asn Leu Asn Gly Val Ile Gln Met Gly Leu Ala Glu Asn Gln Leu Ser
    50                  55                  60

Val Asp Leu Ile Glu Glu Trp Ile Lys Arg Asn Pro Lys Ala Ser Ile
65                  70                  75                  80

Cys Thr Asn Asp Gly Ile Glu Ser Phe Arg Arg Ile Ala Asn Phe Gln
                85                  90                  95

Asp Tyr His Gly Leu Pro Glu Phe Thr Asn Ala Ile Ala Lys Phe Met
            100                 105                 110

Glu Lys Thr Arg Gly Gly Lys Val Lys Phe Asp Ala Lys Arg Val Val
        115                 120                 125

Met Ala Gly Gly Ala Thr Gly Ala Asn Glu Thr Leu Ile Leu Cys Leu
    130                 135                 140

Ala Asp Pro Gly Asp Ala Phe Leu Val Pro Thr Pro Tyr Tyr Pro Gly
145                 150                 155                 160

Phe Asn Arg Asp Leu Arg Trp Arg Ser Gly Val Gln Leu Leu Pro Ile
                165                 170                 175
```

```
Ser Cys Lys Ser Cys Asn Asn Phe Lys Ile Thr Ile Glu Ala Ile Glu
            180                 185                 190

Glu Ala Tyr Glu Lys Gly Gln Gln Ala Asn Val Lys Ile Lys Gly Leu
        195                 200                 205

Ile Leu Thr Asn Pro Cys Asn Pro Leu Gly Thr Ile Leu Asp Arg Asp
    210                 215                 220

Thr Leu Lys Lys Ile Ser Thr Phe Thr Asn Glu His Asn Ile His Leu
225                 230                 235                 240

Val Cys Asp Glu Ile Tyr Ala Ala Thr Val Phe Asn Ser Pro Lys Phe
                245                 250                 255

Val Ser Ile Ala Glu Ile Ile Asn Glu Asp Asn Cys Ile Asn Lys Asp
            260                 265                 270

Leu Val His Ile Val Ser Asn Leu Ser Lys Asp Leu Gly Phe Pro Gly
        275                 280                 285

Phe Arg Val Gly Ile Val Tyr Ser Phe Asn Asp Asp Val Val Asn Cys
    290                 295                 300

Ala Arg Lys Met Ser Ser Phe Gly Leu Val Ser Thr Gln Thr Gln His
305                 310                 315                 320

Leu Leu Ala Phe Met Leu Ser Asp Asp Glu Phe Val Glu Glu Phe Leu
                325                 330                 335

Ile Glu Ser Ala Lys Arg Leu Arg Glu Arg Tyr Glu Lys Phe Thr Arg
            340                 345                 350

Gly Leu Glu Glu Ile Gly Ile Lys Cys Leu Glu Ser Asn Ala Gly Val
        355                 360                 365

Tyr Cys Trp Met Asp Leu Arg Ser Leu Leu Lys Glu Ala Thr Leu Asp
    370                 375                 380

Ala Glu Met Ser Leu Trp Lys Leu Ile Ile Asn Glu Val Lys Leu Asn
385                 390                 395                 400

Val Ser Pro Gly Ser Ser Phe Asn Cys Ser Glu Val Gly Trp Phe Arg
                405                 410                 415

Val Cys Phe Ala Asn Ile Asp Asp Gln Thr Met Glu Ile Ala Leu Ala
```

```
                420                 425                 430

Arg Ile Arg Met Phe Met Asp Ala Tyr Asn Asn Val Asn Lys Asn Gly
        435                 440                 445

Val Met Lys Asn Lys His Asn Gly Arg Gly Thr Thr Tyr Asp Leu Thr
    450                 455                 460

Pro Gln Met Gly Ser Thr Met Lys Met Leu Leu Ala
465                 470                 475
```

<210> 3
<211> 476
<212> PRT
<213> Lycopersicon esculentum

<400> 3

```
Met Asp Leu Glu Thr Ser Glu Ile Ser Asn Tyr Lys Ser Ser Val Val
1               5                  10                  15

Leu Ser Lys Leu Ala Ser Asn Glu Gln His Gly Glu Asn Ser Pro Tyr
            20                  25                  30

Phe Asp Gly Trp Lys Ala Tyr Asp Asn Asp Pro Phe His Leu Val Asn
        35                  40                  45

Asn Leu Asn Gly Val Ile Gln Met Gly Leu Ala Glu Asn Gln Leu Ser
    50                  55                  60

Val Asp Leu Ile Glu Glu Trp Ile Lys Arg Asn Pro Lys Ala Ser Ile
65                  70                  75                  80

Cys Thr Asn Asp Gly Ile Glu Ser Phe Arg Arg Ile Ala Asn Phe Gln
                85                  90                  95

Asp Tyr His Gly Leu Pro Glu Phe Thr Asn Ala Ile Ala Lys Phe Met
            100                 105                 110

Glu Lys Thr Arg Gly Gly Lys Val Lys Phe Asp Ala Lys Arg Val Val
        115                 120                 125

Met Ala Gly Gly Ala Thr Gly Ala Asn Glu Thr Leu Ile Leu Cys Leu
    130                 135                 140

Ala Asp Pro Gly Asp Ala Phe Leu Val Pro Thr Pro Tyr Tyr Pro Gly
145                 150                 155                 160

Phe Asn Arg Asp Leu Arg Trp Arg Ser Gly Val Gln Leu Leu Pro Ile
```

```
                    165                 170                 175

Ser Cys Lys Ser Cys Asn Asn Phe Lys Ile Thr Ile Glu Ala Ile Glu
                180                 185                 190

Glu Ala Tyr Glu Lys Gly Gln Gln Ala Asn Val Lys Ile Lys Gly Leu
        195                 200                 205

Ile Leu Thr Asn Pro Cys Asn Pro Leu Gly Thr Ile Leu Asp Arg Asp
    210                 215                 220

Thr Leu Lys Lys Ile Ser Thr Phe Thr Asn Glu His Asn Ile His Leu
225                 230                 235                 240

Val Cys Asp Glu Ile Tyr Ala Val Thr Val Phe Asn Ser Pro Lys Phe
                245                 250                 255

Val Ser Ile Ala Glu Ile Ile Asn Glu Asp Asn Cys Ile Asn Lys Asp
            260                 265                 270

Leu Val His Ile Val Ser Ser Leu Ser Lys Asp Leu Gly Phe Pro Gly
        275                 280                 285

Phe Arg Val Gly Ile Val Tyr Ser Phe Asn Asp Asp Val Val Asn Cys
    290                 295                 300

Ala Arg Lys Met Ser Ser Phe Gly Leu Val Ser Thr Gln Thr Gln His
305                 310                 315                 320

Leu Leu Ala Phe Met Leu Ser Asp Asp Glu Phe Val Glu Glu Phe Leu
                325                 330                 335

Ile Glu Ser Ala Lys Arg Leu Arg Glu Arg Tyr Glu Lys Phe Thr Arg
            340                 345                 350

Gly Leu Glu Glu Ile Gly Ile Lys Cys Leu Glu Ser Asn Ala Gly Val
        355                 360                 365

Tyr Cys Trp Met Asp Leu Arg Ser Leu Leu Lys Glu Ala Thr Leu Asp
    370                 375                 380

Ala Glu Met Ser Leu Trp Lys Leu Ile Ile Asn Glu Val Lys Leu Asn
385                 390                 395                 400

Val Ser Pro Gly Ser Ser Phe Asn Cys Ser Glu Val Gly Trp Phe Arg
                405                 410                 415
```

```
Val Cys Phe Ala Asn Ile Asp Asp Gln Thr Met Glu Ile Ala Leu Ala
            420                 425                 430

Arg Ile Arg Met Phe Met Asp Ala Tyr Asn Asn Val Asn Lys Asn Gly
        435                 440                 445

Val Met Lys Asn Lys His Asn Gly Arg Gly Thr Thr Tyr Asp Leu Thr
    450                 455                 460

Pro Gln Met Gly Ser Thr Met Lys Met Leu Leu Ala
465                 470                 475
```

<210> 4
<211> 203
<212> PRT
<213> Lycopersicon esculentum

<400> 4

```
Met Asp Leu Glu Thr Ser Glu Ile Ser Asn Tyr Lys Ser Ser Val Val
1               5                   10                  15

Leu Ser Lys Leu Ala Ser Asn Glu Gln His Gly Glu Asn Ser Pro Tyr
            20                  25                  30

Phe Asp Gly Trp Lys Ala Tyr Asp Asn Asp Pro Phe His Leu Val Asn
        35                  40                  45

Asn Leu Asn Gly Val Ile Gln Met Gly Leu Ala Glu Asn Gln Leu Ser
    50                  55                  60

Val Asp Leu Ile Glu Glu Trp Ile Lys Arg Asn Pro Lys Ala Ser Ile
65                  70                  75                  80

Cys Thr Asn Asp Gly Ile Glu Ser Phe Arg Arg Ile Ala Asn Phe Gln
                85                  90                  95

Asp Tyr His Gly Leu Pro Glu Phe Thr Asn Ala Ile Ala Lys Phe Met
            100                 105                 110

Glu Lys Thr Arg Gly Gly Lys Val Lys Phe Asp Ala Lys Arg Val Val
        115                 120                 125

Met Ala Gly Gly Ala Thr Gly Ala Asn Glu Thr Leu Ile Leu Cys Leu
    130                 135                 140

Ala Asp Pro Gly Asp Ala Phe Leu Val Pro Thr Pro Tyr Tyr Pro Gly
145                 150                 155                 160
```

Phe Asn Arg Asp Leu Arg Trp Arg Ser Gly Val Gln Leu Leu Pro Ile
165 170 175

Ser Cys Lys Ser Cys Asn Asn Phe Lys Ile Thr Ile Glu Ala Ile Glu
180 185 190

Glu Ala Tyr Glu Lys Gly Gln Gln Ala Asn Val
195 200

<210> 5
<211> 476
<212> PRT
<213> Lycopersicon esculentum

<400> 5

Met Asp Leu Glu Thr Ser Glu Ile Ser Asn Tyr Lys Ser Ser Val Val
1 5 10 15

Leu Ser Lys Leu Ala Ser Asn Glu Gln His Gly Glu Asn Ser Pro Tyr
20 25 30

Phe Asp Gly Trp Lys Ala Tyr Asp Asn Asp Pro Phe His Leu Val Asn
35 40 45

Asn Leu Asn Gly Val Ile Gln Met Gly Leu Ala Glu Asn Gln Leu Ser
50 55 60

Val Asp Leu Ile Glu Glu Trp Ile Lys Arg Asn Pro Lys Ala Ser Ile
65 70 75 80

Cys Thr Asn Asp Gly Ile Glu Ser Phe Arg Arg Ile Ala Asn Phe Gln
85 90 95

Asp Tyr His Gly Leu Pro Glu Phe Thr Asn Ala Ile Ala Lys Phe Met
100 105 110

Glu Lys Thr Arg Gly Gly Lys Val Lys Phe Asp Ala Lys Arg Val Val
115 120 125

Met Ala Gly Gly Ala Thr Gly Ala Asn Glu Thr Leu Ile Leu Cys Leu
130 135 140

Ala Asp Pro Gly Asp Ala Phe Leu Val Pro Thr Pro Tyr Tyr Pro Gly
145 150 155 160

Phe Asn Arg Asp Leu Arg Trp Arg Ser Gly Val Gln Leu Leu Pro Ile
165 170 175

```
Ser Cys Lys Ser Cys Asn Asn Phe Lys Ile Thr Ile Glu Ala Ile Glu
            180                 185                 190

Glu Ala Tyr Glu Lys Gly Gln Gln Ala Asn Val Lys Ile Lys Gly Leu
        195                 200                 205

Ile Leu Thr Asn Pro Cys Asn Pro Leu Gly Thr Ile Leu Asp Arg Asp
    210                 215                 220

Thr Leu Lys Lys Ile Ser Thr Phe Thr Asn Glu His Asn Ile His Leu
225                 230                 235                 240

Val Cys Asp Glu Ile Tyr Ala Ala Thr Val Phe Asn Ser Pro Lys Phe
                245                 250                 255

Val Ser Ile Ala Glu Ile Ile Asn Glu Asp Asn Cys Ile Asn Lys Asp
            260                 265                 270

Leu Val His Ile Val Ser Ser Leu Ser Lys Asp Leu Gly Phe Pro Gly
        275                 280                 285

Phe Arg Val Gly Ile Val Tyr Ser Phe Asn Asp Asp Val Val Asn Cys
    290                 295                 300

Ala Arg Lys Met Ser Ser Phe Gly Leu Val Ser Thr Gln Thr Gln His
305                 310                 315                 320

Phe Leu Ala Phe Met Leu Ser Asp Asp Glu Phe Val Glu Glu Phe Leu
                325                 330                 335

Ile Glu Ser Ala Lys Arg Leu Arg Glu Arg Tyr Glu Lys Phe Thr Arg
            340                 345                 350

Gly Leu Glu Glu Ile Gly Ile Lys Cys Leu Glu Ser Asn Ala Gly Val
        355                 360                 365

Tyr Cys Trp Met Asp Leu Arg Ser Leu Leu Lys Glu Ala Thr Leu Asp
    370                 375                 380

Ala Glu Met Ser Leu Trp Lys Leu Ile Ile Asn Glu Val Lys Leu Asn
385                 390                 395                 400

Val Ser Pro Gly Ser Ser Phe Asn Cys Ser Glu Val Gly Trp Phe Arg
                405                 410                 415

Val Cys Phe Ala Asn Ile Asp Asp Gln Thr Met Glu Ile Ala Leu Ala
            420                 425                 430
```

```
Arg Ile Arg Met Phe Met Asp Ala Tyr Asn Asn Val Asn Lys Asn Gly
        435                 440                 445

Val Met Lys Asn Lys His Asn Gly Arg Gly Thr Thr Tyr Asp Leu Thr
    450                 455                 460

Pro Gln Met Gly Ser Thr Met Lys Met Leu Leu Ala
465                 470                 475
```

<210> 6
<211> 476
<212> PRT
<213> Lycopersicon esculentum

<400> 6

```
Met Asp Leu Glu Thr Ser Glu Ile Ser Asn Tyr Lys Ser Ser Val Val
1               5                   10                  15

Leu Ser Lys Leu Ala Ser Asn Glu Gln His Gly Glu Asn Ser Pro Tyr
            20                  25                  30

Phe Asp Gly Trp Lys Ala Tyr Asp Asn Asp Pro Phe His Leu Val Asn
        35                  40                  45

Asn Leu Asn Gly Val Ile Gln Met Gly Leu Ala Glu Asn Gln Leu Ser
    50                  55                  60

Val Asp Leu Ile Glu Glu Trp Ile Lys Arg Asn Pro Lys Ala Ser Ile
65                  70                  75                  80

Cys Thr Asn Asp Gly Ile Glu Ser Phe Arg Arg Ile Ala Asn Phe Gln
                85                  90                  95

Asp Tyr His Gly Leu Pro Glu Phe Thr Asn Ala Ile Ala Lys Phe Met
            100                 105                 110

Glu Lys Thr Arg Gly Gly Lys Val Lys Phe Asp Ala Lys Arg Val Val
        115                 120                 125

Met Ala Gly Gly Ala Thr Gly Ala Asn Glu Thr Leu Ile Leu Cys Leu
    130                 135                 140

Ala Asp Pro Gly Asp Ala Phe Leu Val Pro Thr Pro Tyr Tyr Pro Gly
145                 150                 155                 160

Phe Asn Arg Asp Leu Arg Trp Arg Ser Gly Val Gln Leu Leu Pro Ile
                165                 170                 175
```

```
Ser Cys Lys Ser Cys Asn Asn Phe Lys Ile Thr Ile Glu Ala Ile Glu
            180                 185                 190

Glu Ala Tyr Glu Lys Gly Gln Gln Ala Asn Val Lys Ile Lys Gly Leu
        195                 200                 205

Ile Leu Thr Asn Pro Cys Asn Pro Leu Gly Thr Ile Leu Asp Arg Asp
    210                 215                 220

Thr Leu Lys Lys Ile Ser Thr Phe Thr Asn Glu His Asn Ile His Leu
225                 230                 235                 240

Val Cys Asp Glu Ile Tyr Ala Ala Thr Glu Phe Asn Ser Pro Lys Phe
                245                 250                 255

Val Ser Ile Ala Glu Ile Ile Asn Glu Asp Asn Cys Ile Asn Lys Asp
            260                 265                 270

Leu Val His Ile Val Ser Ser Leu Ser Lys Asp Leu Gly Phe Pro Gly
        275                 280                 285

Phe Arg Val Gly Ile Val Tyr Ser Phe Asn Asp Asp Val Val Asn Cys
    290                 295                 300

Ala Arg Lys Met Ser Ser Phe Gly Leu Val Ser Thr Gln Thr Gln His
305                 310                 315                 320

Leu Leu Ala Phe Met Leu Ser Asp Asp Glu Phe Val Glu Glu Phe Leu
                325                 330                 335

Ile Glu Ser Ala Lys Arg Leu Arg Glu Arg Tyr Glu Lys Phe Thr Arg
            340                 345                 350

Gly Leu Glu Glu Ile Gly Ile Lys Cys Leu Glu Ser Asn Ala Gly Val
        355                 360                 365

Tyr Cys Trp Met Asp Leu Arg Ser Leu Leu Lys Glu Ala Thr Leu Asp
    370                 375                 380

Ala Glu Met Ser Leu Trp Lys Leu Ile Ile Asn Glu Val Lys Leu Asn
385                 390                 395                 400

Val Ser Pro Gly Ser Ser Phe Asn Cys Ser Glu Val Gly Trp Phe Arg
                405                 410                 415

Val Cys Phe Ala Asn Ile Asp Asp Gln Thr Met Glu Ile Ala Leu Ala
            420                 425                 430
```

```
Arg Ile Arg Met Phe Met Asp Ala Tyr Asn Asn Val Asn Lys Asn Gly
        435                 440                 445

Val Met Lys Asn Lys His Asn Gly Arg Gly Thr Thr Tyr Asp Leu Thr
    450                 455                 460

Pro Gln Met Gly Ser Thr Met Lys Met Leu Leu Ala
465                 470                 475
```

<210> 7
<211> 476
<212> PRT
<213> Lycopersicon esculentum

<400> 7

```
Met Asp Leu Glu Thr Ser Glu Ile Ser Asn Tyr Lys Ser Ser Val Val
1               5                   10                  15

Leu Ser Lys Leu Ala Ser Asn Glu Gln His Gly Glu Asn Ser Pro Tyr
            20                  25                  30

Phe Asp Gly Trp Lys Ala Tyr Asp Asn Asp Pro Phe His Leu Val Asn
        35                  40                  45

Asn Leu Asn Gly Val Ile Gln Met Gly Leu Ala Glu Asn Gln Leu Ser
    50                  55                  60

Val Asp Leu Ile Glu Glu Trp Ile Lys Arg Asn Pro Lys Ala Ser Ile
65                  70                  75                  80

Cys Thr Asn Asp Gly Ile Glu Ser Phe Arg Arg Ile Ala Asn Phe Gln
                85                  90                  95

Asp Tyr His Gly Leu Pro Glu Phe Thr Asn Ala Ile Ala Lys Phe Met
            100                 105                 110

Glu Lys Thr Arg Gly Gly Lys Val Lys Phe Asp Ala Lys Arg Val Val
        115                 120                 125

Met Ala Gly Gly Ala Thr Gly Ala Asn Glu Thr Leu Ile Leu Cys Leu
    130                 135                 140

Ala Asp Pro Gly Asp Ala Phe Leu Val Pro Thr Pro Tyr Tyr Pro Gly
145                 150                 155                 160

Phe Asn Arg Asp Leu Arg Trp Arg Ser Gly Val Gln Leu Leu Pro Ile
                165                 170                 175
```

```
Ser Cys Lys Ser Cys Asn Asn Phe Lys Ile Thr Ile Glu Ala Ile Glu
            180                 185                 190

Glu Ala Tyr Glu Lys Gly Gln Gln Ala Asn Val Lys Ile Lys Gly Leu
        195                 200                 205

Ile Leu Thr Asn Pro Cys Asn Pro Leu Gly Thr Ile Leu Asp Arg Asp
    210                 215                 220

Thr Leu Lys Lys Ile Ser Thr Phe Thr Asn Glu His Asn Ile His Leu
225                 230                 235                 240

Val Cys Asp Glu Ile Tyr Ala Ala Thr Val Phe Asn Ser Pro Lys Phe
                245                 250                 255

Val Ser Ile Ala Glu Ile Ile Asn Glu Asp Asn Cys Ile Asn Lys Asp
            260                 265                 270

Leu Val His Ile Val Ser Ser Leu Ser Lys Asp Leu Gly Phe Pro Gly
        275                 280                 285

Phe Arg Val Gly Ile Val Tyr Ser Phe Asn Asp Asp Val Val Asn Cys
    290                 295                 300

Ala Arg Lys Met Ser Ser Phe Gly Leu Val Ser Ile Gln Thr Gln His
305                 310                 315                 320

Leu Leu Ala Phe Met Leu Ser Asp Asp Glu Phe Val Glu Glu Phe Leu
                325                 330                 335

Ile Glu Ser Ala Lys Arg Leu Arg Glu Arg Tyr Glu Lys Phe Thr Arg
            340                 345                 350

Gly Leu Glu Glu Ile Gly Ile Lys Cys Leu Glu Ser Asn Ala Gly Val
        355                 360                 365

Tyr Cys Trp Met Asp Leu Arg Ser Leu Leu Lys Glu Ala Thr Leu Asp
    370                 375                 380

Ala Glu Met Ser Leu Trp Lys Leu Ile Ile Asn Glu Val Lys Leu Asn
385                 390                 395                 400

Val Ser Pro Gly Ser Ser Phe Asn Cys Ser Glu Val Gly Trp Phe Arg
                405                 410                 415

Val Cys Phe Ala Asn Ile Asp Asp Gln Thr Met Glu Ile Ala Leu Ala
```

420 425 430

Arg Ile Arg Met Phe Met Asp Ala Tyr Asn Asn Val Asn Lys Asn Gly
435 440 445

Val Met Lys Asn Lys His Asn Gly Arg Gly Thr Thr Tyr Asp Leu Thr
450 455 460

Pro Gln Met Gly Ser Thr Met Lys Met Leu Leu Ala
465 470 475

<210> 8
<211> 1431
<212> DNA
<213> Lycopersicon esculentum

<400> 8

atggatttgg agacgagtga gatttcaaat tacaagtcat cagtagtttt gtctaagttg 60

gctagtaacg aacaacatgg tgaaaactca ccatattttg atgggtggaa agcatacgat 120

aacgatcctt tccacttggt gaataatttg aatggggtta ttcagatggg tctcgcggaa 180

aatcagcttt cagttgactt gattgaagaa tggattaaga gaaatccaaa agcttccatt 240

tgtacaaatg atggaattga atctttcagg agaattgcca actttcaaga ttatcatgga 300

ttgcctgaat tcacaaatgc gattgcaaaa tttatggaga aaacaagagg tggtaaggtt 360

aagtttgatg ctaaacgtgt agtaatggct ggtggagcta ctggagctaa tgagactctc 420

atactttgtt tggctgatcc tggtgatgct tttttagtcc ccacacccta ttacccagga 480

tttaataggg acctaaggtg gagaagtggt gtacaacttt taccaatttc atgcaagagt 540

tgcaataatt tcaaaattac aatagaagct atcgaagagg cctatgaaaa aggtcaacaa 600

gcaaatgtca aaatcaaagg cttgattttg accaaccctt gtaatccatt aggtaccatt 660

ttagataggg acacacttaa aaaaatctcc accttcacta acgaacataa tatccatctt 720

gtttgcgacg aaatatatgc tgctaccgtg ttcaattctc caaaattcgt tagcatcgct 780

gaaattatca acgaagataa ttgtatcaat aaagatttag tacacattgt gtctagtctt 840

tccaaggact taggttttcc aggatttcga gtgggaattg tgtactcatt caacgatgat 900

gttgttaact gtgctagaaa aatgtcgagt tttggtcttg tttcgactca gacacaacat 960

ttgctagctt tcatgttgtc tgacgatgaa tttgtggaag aatttcttat tgaaagcgcg 1020

aaaaggttga gagaaaggta cgagaaattc actagaggac ttgaagaaat aggaatcaag 1080

tgcttagaaa gcaatgcagg ggtttattgt tggatggatt tgcggtcatt gttgaaagaa 1140

gcaacactag atgctgagat gtcactttgg aaactcatca taaacgaagt taagctcaac 1200

gtctcccctg gatcttcgtt caattgctcg gaggtaggat ggtttcgagt ttgttttgca 1260

aatatcgatg atcaaacaat ggagatcgca cttgcaagga ttcggatgtt tatggatgct 1320

tacaacaatg ttaataaaaa tggagtcatg aagaacaagc acaatggaag aggaacaacc 1380

tacgacttaa ctcctcaaat ggggagtacg atgaaaatgt tattagctta a 1431

<210> 9
<211> 1431
<212> DNA
<213> Lycopersicon esculentum

<400> 9

atggatttgg agacgagtga gatttcaaat tacaagtcat cagtagtttt gtctaagttg 60

gctagtaacg aacaacatgg tgaaaactca ccatattttg atgggtggaa agcatacgat 120

aacgatcctt tccacttggt gaataatttg aatggggtta ttcagatggg tctcgcggaa 180

aatcagcttt cagttgactt gattgaagaa tggattaaga gaaatccaaa agcttccatt 240

tgtacaaatg atggaattga atctttcagg agaattgcca actttcaaga ttatcatgga 300

ttgcctgaat tcacaaatgc gattgcaaaa tttatggaga aaacaagagg tggtaaggtt 360

aagtttgatg ctaaacgtgt agtaatggct ggtggagcta ctggagctaa tgagactctc 420

atactttgtt tggctgatcc tggtgatgct tttttagtcc ccacacccta ttacccagga 480

tttaataggg acctaaggtg gagaagtggt gtacaacttt taccaatttc atgcaagagt 540

tgcaataatt tcaaaattac aatagaagct atcgaagagg cctatgaaaa aggtcaacaa 600

gcaaatgtca aaatcaaagg cttgattttg accaaccctt gtaatccatt aggtaccatt 660

ttagataggg acacacttaa aaaaatctcc accttcacta acgaacataa tatccatctt 720

gtttgcgacg aaatatatgc tgctaccgtg ttcaattctc caaaattcgt tagcatcgct 780

gaaattatca acgaagataa ttgtatcaat aaagatttag tacacattgt gtctaatctt 840

tccaaggact taggttttcc aggatttcga gtgggaattg tgtactcatt caacgatgat 900

gttgttaact gtgctagaaa aatgtcgagt tttggtcttg tttcgactca gacacaacat 960

ttgctagctt tcatgttgtc tgacgatgaa tttgtggaag aatttcttat tgaaagcgcg 1020

aaaaggttga gagaaaggta cgagaaattc actagaggac ttgaagaaat aggaatcaag 1080

tgcttagaaa gcaatgcagg ggtttattgt tggatggatt tgcggtcatt gttgaaagaa 1140

gcaacactag atgctgagat gtcactttgg aaactcatca taaacgaagt taagctcaac 1200

gtctcccctg gatcttcgtt caattgctcg gaggtaggat ggtttcgagt ttgttttgca 1260

aatatcgatg atcaaacaat ggagatcgca cttgcaagga ttcggatgtt tatggatgct 1320

tacaacaatg ttaataaaaa tggagtcatg aagaacaagc acaatggaag aggaacaacc 1380

tacgacttaa ctcctcaaat ggggagtacg atgaaaatgt tattagctta a 1431

<210> 10
<211> 1431
<212> DNA
<213> Lycopersicon esculentum

<400> 10

atggatttgg agacgagtga gatttcaaat tacaagtcat cagtagtttt gtctaagttg 60

gctagtaacg aacaacatgg tgaaaactca ccatattttg atgggtggaa agcatacgat 120

aacgatcctt tccacttggt gaataatttg aatggggtta ttcagatggg tctcgcggaa 180

aatcagcttt cagttgactt gattgaagaa tggattaaga gaaatccaaa agcttccatt 240

tgtacaaatg atggaattga atctttcagg agaattgcca actttcaaga ttatcatgga 300

ttgcctgaat tcacaaatgc gattgcaaaa tttatggaga aaacaagagg tggtaaggtt 360

aagtttgatg ctaaacgtgt agtaatggct ggtggagcta ctggagctaa tgagactctc 420

atactttgtt tggctgatcc tggtgatgct tttttagtcc ccacacccta ttacccagga 480

tttaataggg acctaaggtg gagaagtggt gtacaacttt taccaatttc atgcaagagt 540

tgcaataatt tcaaaattac aatagaagct atcgaagagg cctatgaaaa aggtcaacaa 600

gcaaatgtca aaatcaaagg cttgattttg accaaccctt gtaatccatt aggtaccatt 660

ttagataggg acacacttaa aaaaatctcc accttcacta acgaacataa tatccatctt 720

gtttgcgacg aaatatatgc tgttaccgtg ttcaattctc caaaattcgt tagcatcgct 780

gaaattatca acgaagataa ttgtatcaat aaagatttag tacacattgt gtctagtctt 840

tccaaggact taggttttcc aggatttcga gtgggaattg tgtactcatt caacgatgat 900

gttgttaact gtgctagaaa aatgtcgagt tttggtcttg tttcgactca gacacaacat 960

ttgctagctt tcatgttgtc tgacgatgaa tttgtggaag aatttcttat tgaaagcgcg 1020

aaaaggttga gagaaaggta cgagaaattc actagaggac ttgaagaaat aggaatcaag 1080

tgcttagaaa gcaatgcagg ggtttattgt tggatggatt tgcggtcatt gttgaaagaa 1140

gcaacactag atgctgagat gtcactttgg aaactcatca taaacgaagt taagctcaac 1200

gtctcccctg gatcttcgtt caattgctcg gaggtaggat ggtttcgagt ttgttttgca 1260

aatatcgatg atcaaacaat ggagatcgca cttgcaagga ttcggatgtt tatggatgct 1320

tacaacaatg ttaataaaaa tggagtcatg aagaacaagc acaatggaag aggaacaacc 1380

tacgacttaa ctcctcaaat ggggagtacg atgaaaatgt tattagctta a 1431

<210> 11
<211> 1431
<212> DNA
<213> Lycopersicon esculentum

<400> 11

atggatttgg agacgagtga gatttcaaat tacaagtcat cagtagtttt gtctaagttg 60

gctagtaacg aacaacatgg tgaaaactca ccatattttg atgggtggaa agcatacgat 120

aacgatcctt tccacttggt gaataatttg aatggggtta ttcagatggg tctcgcggaa 180

aatcagcttt cagttgactt gattgaagaa tggattaaga gaaatccaaa agcttccatt 240

tgtacaaatg atggaattga atctttcagg agaattgcca actttcaaga ttatcatgga 300

ttgcctgaat tcacaaatgc gattgcaaaa tttatggaga aaacaagagg tggtaaggtt 360

aagtttgatg ctaaacgtgt agtaatggct ggtggagcta ctggagctaa tgagactctc 420

atactttgtt tggctgatcc tggtgatgct tttttagtcc ccacacccta ttacccagga 480

tttaataggg acctaaggtg gagaagtggt gtacaacttt taccaatttc atgcaagagt 540

tgcaataatt tcaaaattac aatagaagct atcgaagagg cctatgaaaa aggtcaacaa 600

gcaaatgtct aaatcaaagg cttgattttg accaaccctt gtaatccatt aggtaccatt 660

ttagataggg acacacttaa aaaaatctcc accttcacta acgaacataa tatccatctt 720

gtttgcgacg aaatatatgc tgctaccgtg ttcaattctc caaaattcgt tagcatcgct 780

gaaattatca acgaagataa ttgtatcaat aaagatttag tacacattgt gtctagtctt 840

tccaaggact taggttttcc aggatttcga gtgggaattg tgtactcatt caacgatgat 900

gttgttaact gtgctagaaa aatgtcgagt tttggtcttg tttcgactca gacacaacat 960

ttgctagctt tcatgttgtc tgacgatgaa tttgtggaag aatttcttat tgaaagcgcg 1020

aaaaggttga gagaaaggta cgagaaattc actagaggac ttgaagaaat aggaatcaag 1080

tgcttagaaa gcaatgcagg ggtttattgt tggatggatt tgcggtcatt gttgaaagaa 1140

gcaacactag atgctgagat gtcactttgg aaactcatca taaacgaagt taagctcaac 1200

gtctcccctg gatcttcgtt caattgctcg gaggtaggat ggtttcgagt ttgttttgca 1260

aatatcgatg atcaaacaat ggagatcgca cttgcaagga ttcggatgtt tatggatgct 1320

tacaacaatg ttaataaaaa tggagtcatg aagaacaagc acaatggaag aggaacaacc 1380

tacgacttaa ctcctcaaat ggggagtacg atgaaaatgt tattagctta a 1431

<210> 12
<211> 1431
<212> DNA
<213> Lycopersicon esculentum

<400> 12

atggatttgg agacgagtga gatttcaaat tacaagtcat cagtagtttt gtctaagttg 60

gctagtaacg aacaacatgg tgaaaactca ccatattttg atgggtggaa agcatacgat 120

aacgatcctt tccacttggt gaataatttg aatggggtta ttcagatggg tctcgcggaa 180

aatcagcttt cagttgactt gattgaagaa tggattaaga gaaatccaaa agcttccatt 240

tgtacaaatg atggaattga atctttcagg agaattgcca actttcaaga ttatcatgga 300

ttgcctgaat tcacaaatgc gattgcaaaa tttatggaga aaacaagagg tggtaaggtt 360

aagtttgatg ctaaacgtgt agtaatggct ggtggagcta ctggagctaa tgagactctc 420

atactttgtt tggctgatcc tggtgatgct tttttagtcc ccacacccta ttacccagga 480

tttaataggg acctaaggtg gagaagtggt gtacaacttt taccaatttc atgcaagagt 540

tgcaataatt tcaaaattac aatagaagct atcgaagagg cctatgaaaa aggtcaacaa 600

gcaaatgtca aaatcaaagg cttgattttg accaaccctt gtaatccatt aggtaccatt 660

ttagataggg acacacttaa aaaaatctcc accttcacta acgaacataa tatccatctt 720

gtttgcgacg aaatatatgc tgctaccgtg ttcaattctc caaaattcgt tagcatcgct 780

gaaattatca acgaagataa ttgtatcaat aaagatttag tacacattgt gtctagtctt 840

tccaaggact taggttttcc aggatttcga gtgggaattg tgtactcatt caacgatgat 900

gttgttaact gtgctagaaa aatgtcgagt tttggtcttg tttcgactca gacacaacat 960

tttctagctt tcatgttgtc tgacgatgaa tttgtggaag aatttcttat tgaaagcgcg 1020

aaaaggttga gagaaaggta cgagaaattc actagaggac ttgaagaaat aggaatcaag 1080

tgcttagaaa gcaatgcagg ggtttattgt tggatggatt tgcggtcatt gttgaaagaa 1140

gcaacactag atgctgagat gtcactttgg aaactcatca taaacgaagt taagctcaac 1200

gtctcccctg gatcttcgtt caattgctcg gaggtaggat ggtttcgagt ttgttttgca 1260

aatatcgatg atcaaacaat ggagatcgca cttgcaagga ttcggatgtt tatggatgct 1320

tacaacaatg ttaataaaaa tggagtcatg aagaacaagc acaatggaag aggaacaacc 1380

tacgacttaa ctcctcaaat ggggagtacg atgaaaatgt tattagctta a 1431

<210> 13
<211> 1431
<212> DNA
<213> Lycopersicon esculentum

<400> 13

atggatttgg agacgagtga gatttcaaat tacaagtcat cagtagtttt gtctaagttg 60

gctagtaacg aacaacatgg tgaaaactca ccatattttg atgggtggaa agcatacgat 120

aacgatcctt tccacttggt gaataatttg aatggggtta ttcagatggg tctcgcggaa 180

aatcagcttt cagttgactt gattgaagaa tggattaaga gaaatccaaa agcttccatt 240

tgtacaaatg atggaattga atctttcagg agaattgcca actttcaaga ttatcatgga 300

ttgcctgaat tcacaaatgc gattgcaaaa tttatggaga aaacaagagg tggtaaggtt 360

aagtttgatg ctaaacgtgt agtaatggct ggtggagcta ctggagctaa tgagactctc 420

atactttgtt tggctgatcc tggtgatgct tttttagtcc ccacacccta ttacccagga 480

tttaataggg acctaaggtg gagaagtggt gtacaacttt taccaatttc atgcaagagt 540

tgcaataatt tcaaaattac aatagaagct atcgaagagg cctatgaaaa aggtcaacaa 600

gcaaatgtca aaatcaaagg cttgattttg accaaccctt gtaatccatt aggtaccatt 660

ttagataggg acacacttaa aaaaatctcc accttcacta acgaacataa tatccatctt 720

gtttgcgacg aaatatatgc tgctaccgag ttcaattctc caaaattcgt tagcatcgct 780

gaaattatca acgaagataa ttgtatcaat aaagatttag tacacattgt gtctagtctt 840

tccaaggact taggttttcc aggatttcga gtgggaattg tgtactcatt caacgatgat 900

gttgttaact gtgctagaaa aatgtcgagt tttggtcttg tttcgactca gacacaacat 960

ttgctagctt tcatgttgtc tgacgatgaa tttgtggaag aatttcttat tgaaagcgcg 1020

aaaaggttga gagaaaggta cgagaaattc actagaggac ttgaagaaat aggaatcaag 1080

tgcttagaaa gcaatgcagg ggtttattgt tggatggatt tgcggtcatt gttgaaagaa 1140

gcaacactag atgctgagat gtcactttgg aaactcatca taaacgaagt taagctcaac 1200

gtctcccctg gatcttcgtt caattgctcg gaggtaggat ggtttcgagt ttgttttgca 1260

aatatcgatg atcaaacaat ggagatcgca cttgcaagga ttcggatgtt tatggatgct 1320

tacaacaatg ttaataaaaa tggagtcatg aagaacaagc acaatggaag aggaacaacc 1380

tacgacttaa ctcctcaaat ggggagtacg atgaaaatgt tattagctta a 1431

<210> 14
<211> 1431
<212> DNA
<213> Lycopersicon esculentum

<400> 14

atggatttgg agacgagtga gatttcaaat tacaagtcat cagtagtttt gtctaagttg 60

gctagtaacg aacaacatgg tgaaaactca ccatattttg atgggtggaa agcatacgat 120

aacgatcctt tccacttggt gaataatttg aatggggtta ttcagatggg tctcgcggaa 180

aatcagcttt cagttgactt gattgaagaa tggattaaga gaaatccaaa agcttccatt 240

tgtacaaatg atggaattga atctttcagg agaattgcca actttcaaga ttatcatgga 300

ttgcctgaat tcacaaatgc gattgcaaaa tttatggaga aaacaagagg tggtaaggtt 360

aagtttgatg ctaaacgtgt agtaatggct ggtggagcta ctggagctaa tgagactctc 420

atactttgtt tggctgatcc tggtgatgct tttttagtcc ccacacccta ttacccagga 480

tttaataggg acctaaggtg gagaagtggt gtacaacttt taccaatttc atgcaagagt 540

tgcaataatt tcaaaattac aatagaagct atcgaagagg cctatgaaaa aggtcaacaa 600

gcaaatgtca aaatcaaagg cttgattttg accaaccctt gtaatccatt aggtaccatt 660

ttagataggg acacacttaa aaaaatctcc accttcacta acgaacataa tatccatctt 720

gtttgcgacg aaatatatgc tgctaccgtg ttcaattctc caaaattcgt tagcatcgct 780

gaaattatca acgaagataa ttgtatcaat aaagatttag tacacattgt gtctagtctt 840

tccaaggact taggttttcc aggatttcga gtgggaattg tgtactcatt caacgatgat 900

gttgttaact gtgctagaaa aatgtcgagt tttggtcttg tttcgattca gacacaacat 960

ttgctagctt tcatgttgtc tgacgatgaa tttgtggaag aatttcttat tgaaagcgcg 1020

aaaaggttga gagaaaggta cgagaaattc actagaggac ttgaagaaat aggaatcaag 1080

tgcttagaaa gcaatgcagg ggtttattgt tggatggatt tgcggtcatt gttgaaagaa 1140

gcaacactag atgctgagat gtcactttgg aaactcatca taaacgaagt taagctcaac 1200

gtctcccctg gatcttcgtt caattgctcg gaggtaggat ggtttcgagt ttgttttgca 1260

aatatcgatg atcaaacaat ggagatcgca cttgcaagga ttcggatgtt tatggatgct 1320

tacaacaatg ttaataaaaa tggagtcatg aagaacaagc acaatggaag aggaacaacc 1380

tacgacttaa ctcctcaaat ggggagtacg atgaaaatgt tattagctta a 1431

<210> 15
<211> 2641
<212> DNA
<213> Lycopersicon esculentum

<220>
<221> exon
<222> (1)..(318)
<223> exon1

<220>
<221> exon

<222> (796)..(955)
<223> exon2

<220>
<221> exon
<222> (1689)..(2641)
<223> exon2

<400> 15

```
atg gat ttg gag acg agt gag att tca aat tac aag tca tca gta gtt          48
Met Asp Leu Glu Thr Ser Glu Ile Ser Asn Tyr Lys Ser Ser Val Val
1               5                   10                  15

ttg tct aag ttg gct agt aac gaa caa cat ggt gaa aac tca cca tat          96
Leu Ser Lys Leu Ala Ser Asn Glu Gln His Gly Glu Asn Ser Pro Tyr
            20                  25                  30

ttt gat ggg tgg aaa gca tac gat aac gat cct ttc cac ttg gtg aat         144
Phe Asp Gly Trp Lys Ala Tyr Asp Asn Asp Pro Phe His Leu Val Asn
        35                  40                  45

aat ttg aat ggg gtt att cag atg ggt ctc gcg gaa aat cag ctt tca         192
Asn Leu Asn Gly Val Ile Gln Met Gly Leu Ala Glu Asn Gln Leu Ser
    50                  55                  60

gtt gac ttg att gaa gaa tgg att aag aga aat cca aaa gct tcc att         240
Val Asp Leu Ile Glu Glu Trp Ile Lys Arg Asn Pro Lys Ala Ser Ile
65                  70                  75                  80
```

```
tgt aca aat gat gga att gaa tct ttc agg aga att gcc aac ttt caa        288
Cys Thr Asn Asp Gly Ile Glu Ser Phe Arg Arg Ile Ala Asn Phe Gln
                 85                  90                  95

gat tat cat gga ttg cct gaa ttc aca aat gtaagttttg ttatttctct        338
Asp Tyr His Gly Leu Pro Glu Phe Thr Asn
            100                 105

cctttcaaaa acaaaatgtc acattaaaaa ttagtatatt tttttagtta tcctccgttc    398

aattcttaag aaatatctaa taaataaaag gattattttt cttaataggc gtgaaataaa    458

ttaaacttag acttctttta agatgaatat gaaatactta ctactattat atatgaattg    518

tagcggtgaa agtcattata aatttgtaca aaaaaaaaaa agaaagttaa tgatcaattt    578

tattactata attttacatt tacttggaat aaagaactaa gattacattt agttgaaata    638

gatatatttc ttgacttcta tctcatacat atatttcatt ttatctgaca ctatttttac    698

ttgtttattg aaaatttaaa aaattacata cgttattaaa ataaatatat ttttatccaa    758

ttttttcgta taaaaaaata ttttttttgt tgtgtag gcg att gca aaa ttt atg      813
                                        Ala Ile Ala Lys Phe Met
                                                    110

gag aaa aca aga ggt ggt aag gtt aag ttt gat gct aaa cgt gta gta       861
Glu Lys Thr Arg Gly Gly Lys Val Lys Phe Asp Ala Lys Arg Val Val
        115                 120                 125

atg gct ggt gga gct act gga gct aat gag act ctc ata ctt tgt ttg       909
Met Ala Gly Gly Ala Thr Gly Ala Asn Glu Thr Leu Ile Leu Cys Leu
    130                 135                 140

gct gat cct ggt gat gct ttt tta gtc ccc aca ccc tat tac cca g         955
Ala Asp Pro Gly Asp Ala Phe Leu Val Pro Thr Pro Tyr Tyr Pro
145                 150                 155

ggtatgtata catatttcta aattgaattc aacttatatt atatcgatcg tgaaaaaaat    1015

agatgtttct taattaaata ttaaattcct tcttgcctat ttaaaatgcg aattatatta    1075

tatcactcag ttgaccttta aaatcgatat tatataatat aatttgtatt tttcaaatta    1135

aataagtgaa aatagtcatt taatttattt agtagataag taaaaatgga cggacgaagt    1195

atataaatac catttcaggt aattgattga ggggagattt ttttttaatg aaataacact    1255

ttagtaattt aagagagact atccgtattg gtcaaatctt tagaactaat attaatgggg    1315

gcaacttttc ctacttgtga tttgtcaaaa aagttgaaaa ctaacaaaca aatgatattt    1375

ttggttctct gtctttttat tttgattaaa aaaaatgaaa ttttctcatt ttttaaaaaa    1435

aataatggat agtataaaaa ttataattac tttctttatc ctgatttatg taacatagtt    1495

tgaatttcta gacgtttcaa gtttaatttt gaatttgtat catataaaaa gtatttattt    1555

attttaaaat cgttatgatt aataattcaa atgaaaaaat tgattgactc tcgaaatttg    1615

tgattataca atgataaaat tatttaatgg cttacattgg catgtattta agtcttcatt    1675
```

```
tgtattaaat gca ga  ttt aat agg gac cta agg tgg aga agt ggt gta        1723
               Gly Phe Asn Arg Asp Leu Arg Trp Arg Ser Gly Val
               160                 165                 170

caa ctt tta cca att tca tgc aag agt tgc aat aat ttc aaa att aca        1771
Gln Leu Leu Pro Ile Ser Cys Lys Ser Cys Asn Asn Phe Lys Ile Thr
            175                 180                 185

ata gaa gct atc gaa gag gcc tat gaa aaa ggt caa caa gca aat gtc        1819
Ile Glu Ala Ile Glu Glu Ala Tyr Glu Lys Gly Gln Gln Ala Asn Val
        190                 195                 200

aaa atc aaa ggc ttg att ttg acc aac cct tgt aat cca tta ggt acc        1867
Lys Ile Lys Gly Leu Ile Leu Thr Asn Pro Cys Asn Pro Leu Gly Thr
    205                 210                 215

att tta gat agg gac aca ctt aaa aaa atc tcc acc ttc act aac gaa        1915
Ile Leu Asp Arg Asp Thr Leu Lys Lys Ile Ser Thr Phe Thr Asn Glu
220                 225                 230                 235

cat aat atc cat ctt gtt tgc gac gaa ata tat gct gct acc gtg ttc        1963
His Asn Ile His Leu Val Cys Asp Glu Ile Tyr Ala Ala Thr Val Phe
                240                 245                 250

aat tct cca aaa ttc gtt agc atc gct gaa att atc aac gaa gat aat        2011
Asn Ser Pro Lys Phe Val Ser Ile Ala Glu Ile Ile Asn Glu Asp Asn
            255                 260                 265

tgt atc aat aaa gat tta gta cac att gtg tct agt ctt tcc aag gac        2059
Cys Ile Asn Lys Asp Leu Val His Ile Val Ser Ser Leu Ser Lys Asp
        270                 275                 280

tta ggt ttt cca gga ttt cga gtg gga att gtg tac tca ttc aac gat        2107
Leu Gly Phe Pro Gly Phe Arg Val Gly Ile Val Tyr Ser Phe Asn Asp
    285                 290                 295

gat gtt gtt aac tgt gct aga aaa atg tcg agt ttt ggt ctt gtt tcg        2155
Asp Val Val Asn Cys Ala Arg Lys Met Ser Ser Phe Gly Leu Val Ser
300                 305                 310                 315

act cag aca caa cat ttg cta gct ttc atg ttg tct gac gat gaa ttt        2203
Thr Gln Thr Gln His Leu Leu Ala Phe Met Leu Ser Asp Asp Glu Phe
                320                 325                 330

gtg gaa gaa ttt ctt att gaa agc gcg aaa agg ttg aga gaa agg tac        2251
Val Glu Glu Phe Leu Ile Glu Ser Ala Lys Arg Leu Arg Glu Arg Tyr
            335                 340                 345

gag aaa ttc act aga gga ctt gaa gaa ata gga atc aag tgc tta gaa        2299
Glu Lys Phe Thr Arg Gly Leu Glu Glu Ile Gly Ile Lys Cys Leu Glu
        350                 355                 360

agc aat gca ggg gtt tat tgt tgg atg gat ttg cgg tca ttg ttg aaa        2347
Ser Asn Ala Gly Val Tyr Cys Trp Met Asp Leu Arg Ser Leu Leu Lys
    365                 370                 375

gaa gca aca cta gat gct gag atg tca ctt tgg aaa ctc atc ata aac        2395
Glu Ala Thr Leu Asp Ala Glu Met Ser Leu Trp Lys Leu Ile Ile Asn
380                 385                 390                 395

gaa gtt aag ctc aac gtc tcc cct gga tct tcg ttc aat tgc tcg gag        2443
Glu Val Lys Leu Asn Val Ser Pro Gly Ser Ser Phe Asn Cys Ser Glu
                400                 405                 410
```

```
gta gga tgg ttt cga gtt tgt ttt gca aat atc gat gat caa aca atg       2491
Val Gly Trp Phe Arg Val Cys Phe Ala Asn Ile Asp Asp Gln Thr Met
            415                 420                 425

gag atc gca ctt gca agg att cgg atg ttt atg gat gct tac aac aat       2539
Glu Ile Ala Leu Ala Arg Ile Arg Met Phe Met Asp Ala Tyr Asn Asn
        430                 435                 440

gtt aat aaa aat gga gtc atg aag aac aag cac aat gga aga gga aca       2587
Val Asn Lys Asn Gly Val Met Lys Asn Lys His Asn Gly Arg Gly Thr
    445                 450                 455

acc tac gac tta act cct caa atg ggg agt acg atg aaa atg tta tta       2635
Thr Tyr Asp Leu Thr Pro Gln Met Gly Ser Thr Met Lys Met Leu Leu
460                 465                 470                 475

gct taa                                                              2641
Ala
```

Claims

1. A cultivated plant of the species *Solanum lycopersicum* comprising an *acs* 4 allele having one or more mutations, said mutations resulting in production of a mutant acs 4 protein having loss-of-function or reduced function compared to wild type Acs 4 protein of SEQ ID NO: 1 wherein said mutation or mutations result in reduced ethylene production and/or delayed fruit ripening and/or a longer shelf life compared to *Solanum lycopersicum* being homozygous for the wild type *ACS 4* allele.

2. The cultivated plant according to claim 1 wherein said mutation or mutations result in the tomato fruits requiring significantly more days to reach the red stage compared to *Solanum lycopersicum* being homozygous for the wild type Acs 4 allele.

3. The cultivated plant according to anyone of claims 1 to 2 wherein said mutation or mutations result in the tomato fruits of said plant having at least an 20 % reduced ethylene production compared to *Solanum lycopersicum* being homozygous for the wild type *Acs4* allele.

4. The plant according to any one of claims 1 to 3 wherein the loss-of-function or reduced function of the mutant acs 4 protein is due to one or more amino acids being deleted, replaced and/or inserted compared to the wild type Acs 4 protein of SEQ. ID NO: 1.

5. The plant according to any one of the preceding claims, wherein said mutant acs4 protein has a functional large domain.

6. The plant according to any one of claims 1 to 4, wherein said mutant acs4 protein has functional small domains.

7. The plant according to any one of the preceding claims, wherein said loss-of-function or reduced function of the mutant acs4 protein is due to one or more amino acids being deleted, replaced and/or inserted in the large domain.

8. The plant according to any one of the preceding claims, wherein said mutant acs 4 protein has one or more amino acids changed selected from the group consisting of A248V, S279N, L321F, V250E, S253P, and T316I; or wherein in said mutant acs4 protein misses all amino acids 204 to 476 of SEQ ID NO: 1.

9. The plant according to claims 1 to 8, wherein the plant is an F1 hybrid plant.

10. Seeds from which a plant according to any one of the preceding claims can be grown.

11. Tomato fruit, seeds, pollen, or progeny of the plant of anyone of claims 1-9 comprising an *acs4* allele having one or more mutations, said mutations resulting in the production of a mutant acs4 protein having loss-of-function or reduced function compared to wild type Acs4 protein.

12. The tomato fruit of claim 11, wherein the tomato fruit has reduced ethylene production and/or delayed ripening and/or an increased shelf life compared to fruits from *Solanum lycopersicum* plants being homozygous for the wild type *Acs4* allele.

13. The fruit according to claim 12, wherein the shelf life is at least 2 days longer than the shelf life of a tomato fruit being homozygous for the wild type *Acs* 4 allele.

14. The fruit according to claim 12, wherein the reduced ethylene production is at least 15 % reduced compared to *Solanum lycopersicum* being homozygous for the wild type *Acs4* allele.

15. The plant according to claim 1, wherein said mutant acs 4 protein has one or more amino acids changed selected from the group consisting of A248, S279, L321, V250, and T316; or
wherein the *acs* 4 allele is the *acs* 4 allele as found in seed deposited under accession number NCIMB 42034, NCIMB 42037, NCIMB 42038, NCIMB 42039, or NCIMB 42041 in one or two copies.

Patentansprüche

1. Kultivierte Pflanze der Spezies *Solanum lycopersicum,* umfassend ein *acs* 4-Allel mit einer oder mehreren Mutationen, wobei die Mutationen zur Produktion eines mutierten acs 4-Proteins mit Funktionsverlust (Loss-of-function) oder Funktionsminderung verglichen mit Wildtyp-Acs-4-Protein unter SEQ ID NO: 1 führen, wobei die Mutation bzw. Mutationen zu verminderter Ethylenproduktion und/oder verzögerter Fruchtreifung und/oder längerer Haltbarkeit verglichen mit *Solanum lycopersicum,* die für das Wildtyp-ACS 4-Allel homozygot ist, führt bzw. führen.

2. Kultivierte Pflanze nach Anspruch 1, wobei die Mutation oder Mutationen dazu führen, dass, verglichen mit *Solanum lycopersicum,* die für das Wildtyp-Acs-4-Allel homozygot ist, die Tomatenfrüchte deutlich mehr Tage bis zum Erreichen des Rotstadiums benötigen.

3. Kultivierte Pflanze nach einem der Ansprüche 1 bis 2, wobei die Mutation oder Mutationen dazu führen, dass, verglichen mit *Solanum lycopersicum,* die für das Wildtyp-Acs4-Allel homozygot ist, die Tomatenfrüchte der Pflanze eine um wenigstens 20% verminderte Ethylenproduktion aufweisen.

4. Pflanze nach einem der Ansprüche 1 bis 3, wobei der Funktionsverlust bzw. die Funktionsminderung des mutierten acs 4-Proteins auf die Deletion, Ersetzung und/oder Insertion einer oder mehrerer Aminosäuren gegenüber dem Wildtyp-Acs4-Protein unter SEQ. ID NO: 1 zurückzuführen ist.

5. Pflanze nach einem der vorhergehenden Ansprüche, wobei das mutierte acs4-Protein eine funktionsfähige große Domäne aufweist.

6. Pflanze nach einem der Ansprüche 1 bis 4, wobei das mutierte acs4-Protein funktionsfähige kleine Domänen aufweist.

7. Pflanze nach einem der vorhergehenden Ansprüche, wobei der Funktionsverlust bzw. die Funktionsminderung des mutierten acs4-Proteins auf die Deletion, Ersetzung und/oder Insertion einer oder mehrerer Aminosäuren in der großen Domäne zurückzuführen ist.

8. Pflanze nach einem der vorhergehenden Ansprüche, wobei beim mutierten acs 4-Protein Änderungen einer oder mehrerer Aminosäuren vorliegen, die ausgewählt sind aus der Gruppe bestehend aus A248V, S279N, L321F, V250E, S253P und T316I; oder wobei beim mutierten acs4-Protein alle Aminosäuren 204 bis 476 von SEQ ID NO: 1 fehlen.

9. Pflanze nach Anspruch 1 bis 8, wobei es sich bei der Pflanze um eine F1-Hybride handelt.

10. Samen, aus denen eine Pflanze nach einem der vorhergehenden Ansprüche gezogen werden kann.

11. Tomatenfrucht, Samen, Pollen oder Nachkommen der Pflanze nach einem der Ansprüche 1-9, umfassend ein *acs4*-Allel mit einer oder mehreren Mutationen, wobei die Mutationen zur Produktion eines mutierten acs4-Proteins mit Funktionsverlust oder Funktionsminderung verglichen mit Wildtyp-Acs4-Protein führen.

12. Tomatenfrucht nach Anspruch 11, wobei die Tomatenfrucht verminderte Ethylenproduktion und/oder verzögerte Fruchtreifung und/oder eine längere Haltbarkeit verglichen mit Früchten von *Solanum lycopersicum*-Pflanzen, die für das Wildtyp-Acs4-Allel homozygot sind, aufweist.

13. Frucht nach Anspruch 12, wobei die Haltbarkeit wenigstens 2 Tage länger ist als die Haltbarkeit einer Tomatenfrucht, die für das Wildtyp-Acs 4-Allel homozygot ist.

14. Frucht nach Anspruch 12, wobei die verminderte Ethylenproduktion um wenigstens 15 % verglichen mit *Solanum lycopersicum,* die für das Wildtyp-*Acs4*-Allel homozygot ist, vermindert ist.

15. Pflanze nach Anspruch 1, wobei beim mutierten acs 4-Protein eine oder mehrere Aminosäuren ausgewählt aus der Gruppe bestehend aus A248, S279, L321, V250 und T316 geändert sind; oder
wobei es sich bei dem *acs 4*-Allel um das *acs 4*-Allel handelt, wie es in Samen hinterlegt unter Zugangsnummer NCIMB 42034, NCIMB 42037, NCIMB 42038, NCIMB 42039 oder NCIMB 42041 in einer oder zwei Kopien vorliegt.

Revendications

1. Plante cultivée de l'espèce *Solanum lycopersicum* comprenant un allèle *acs 4* ayant une ou plusieurs mutation(s), lesdites mutations conduisant à la production d'une protéine acs 4 mutante ayant une perte de fonction ou une fonction réduite en comparaison avec une protéine Acs 4 de type sauvage de SEQ ID n° : 1 dans laquelle ladite mutation ou lesdites mutations conduit/conduisent à une production réduite d'éthylène et/ou à un mûrissement retardé du fruit et/ou à une durée de conservation allongée en comparaison avec une *Solanum lycopersicum* étant homozygote pour l'allèle *ACS* 4 de type sauvage.

2. Plante cultivée selon la revendication 1, dans laquelle ladite mutation ou lesdites mutations conduit/conduisent au fait que les tomates-fruits requièrent significativement plus de jours pour atteindre le stade rouge en comparaison avec une *Solanum lycopersicum* étant homozygote pour l'allèle Acs 4 de type sauvage.

3. Plante cultivée selon l'une quelconque des revendications 1 et 2, dans laquelle ladite mutation ou lesdites mutations conduit/conduisent au fait que les tomates-fruits de ladite plante aient une production d'éthylène réduite d'au moins 20 % en comparaison avec une *Solanum lycopersicum* étant homozygote pour l'allèle *Acs4* de type sauvage.

4. Plante selon l'une quelconque des revendications 1 à 3, dans laquelle la perte de fonction ou la fonction réduite de la protéine acs 4 mutante est due à une délétion, à un remplacement et/ou à une insertion d'un ou de plusieurs acide(s) aminé(s) en comparaison avec la protéine Acs 4 de type sauvage de SEQ ID n° : 1.

5. Plante selon l'une quelconque des revendications précédentes, dans laquelle ladite protéine acs4 mutante a un grand domaine fonctionnel.

6. Plante selon l'une quelconque des revendications 1 à 4, dans laquelle ladite protéine acs4 mutante a de petits domaines fonctionnels.

7. Plante selon l'une quelconque des revendications précédentes, dans laquelle ladite perte de fonction ou ladite fonction réduite de la protéine acs4 mutante est due à une délétion, à un remplacement et/ou à une insertion d'un ou de plusieurs acide(s) aminé(s) dans le grand domaine.

8. Plante selon l'une quelconque des revendications précédentes, dans laquelle ladite protéine acs 4 mutante a un ou plusieurs acide(s) aminé(s) changé(s) choisi(s) dans le groupe constitué par les A248V, S279N, L321F, V250E, S253P et T316I ; ou dans laquelle il manque dans ladite protéine acs4 mutante tous les acides aminés 204 à 476 de la SEQ ID n° : 1.

9. Plante selon les revendications 1 à 8, la plante étant une plante hybride F1.

10. Graines à partir desquelles peut être développée une plante selon l'une quelconque des revendications précédentes.

11. Tomate-fruit, graines, pollen ou descendance de la plante de l'une quelconque des revendications 1-9 comprenant un allèle *acs4* ayant une ou plusieurs mutation(s), lesdites mutations conduisant à la production d'une protéine acs4

mutante ayant une perte de fonction ou une fonction réduite en comparaison avec une protéine Acs4 de type sauvage.

12. Tomate-fruit de la revendication 11, la tomate-fruit ayant une production réduite d'éthylène et/ou un mûrissement retardé et/ou une durée de conservation augmentée en comparaison avec des fruits provenant de plantes *Solanum lycopersicum* étant homozygotes pour l'allèle *Acs4* de type sauvage.

13. Fruit selon la revendication 12, dans lequel la durée de conservation est plus longue d'au moins 2 jours que la durée de conservation d'une tomate-fruit étant homozygote pour l'allèle *Acs 4* de type sauvage.

14. Fruit selon la revendication 12, dans lequel la production réduite d'éthylène est réduite d'au moins 15 % en comparaison avec une *Solanum lycopersicum* étant homozygote pour l'allèle *Acs4* de type sauvage.

15. Plante selon la revendication 1, dans laquelle ladite protéine acs 4 mutante a un ou plusieurs acide(s) aminé(s) changé(s) choisi(s) dans le groupe constitué par les A248, S279, L321, V250 et T316 ; ou
dans laquelle l'allèle *acs 4* est l'allèle *acs 4* tel qu'on le trouve dans la graine ayant fait l'objet d'un dépôt sous les numéros d'accession NCIMB 42034, NCIMB 42037, NCIMB 42038, NCIMB 42039 ou NCIMB 42041 dans une ou deux copie(s).

Figure 1



Q42668 MGFHQTH--- ETNQVLLSKI AISDGHGEDS PYFDGWKAYD NDPFHPENNP LGVIQMGLAE NQLSPDLIVD 67
Q43810 M-----E--- NKSKQLLSKI ATNDGHGENS PYFDGWKAYD RDPFHPSQNP NGVIQMGLAE NQLSSDLIED 62
Q96579 MVSISKN--- NQKQQLLSKI ATNDGHGENS PYFDGWKAYA NNPFHLTDNP TGVIQMGLAE NQLCFDLIQE 67
Q43747 MVALTAE--- KQDQNLLSRM AAGDGHGEKS AYFDGWKAYE ENPFHPIDRP DGVIQMGLAE NQLCGDLMRK 67
Le-ACS4 MDLETSEISN YKSSVVLSKL ASNEQHGENS PYFDGWKAYD NDPFHLVNNL NGVIQMGLAE NQLSVDLIEE 70
Q41688 M--------- ----RLLSTI ATCNSHGQDS SYFLGWQEYE KNPYDELHNP KGIIQMGLAE NQLSFDLLES 57
Q43166 M--------- ----KLLSKK AMCNSHGQDS SYFLGWEEYE KNPYDETRNP KGIIQMGLAE NQLSFDLLES 57
Conservation
Q42668 WIRKHPKASI YTDEGLLNFK DIANFQDYHG LPEFRNAIAS FMGKARGGRV KFDPSRIVMG GGATGASETV 137
Q43810 WVRSNPEASI CTLEGVGKFK DVANFQDYHG LLEFRHAVAK FMSRGRGGKV TFDPDRVVMS GGATGANELI 132
Q96579 WVVNNPKASI CTVEGAENFQ DIAIFQDYHG LPEFRQAVAR FMEKVRGDRV TFDPNRIVMS GGATGAHEML 137
Q43747 WVLEHPEASI CTAEGVNQFS DIAIFQDYHG LPEFRQAVAK FMEKTRNNKV KFDPDRIVMS GGATGAHETV 137
Le-ACS4 WIKRNPKASI CTNDGIESFR RIANFQDYHG LPEFTNAIAK FMEKTRGGKV KFDAKRVVMA GGATGANETL 140
Q41688 WLAKNPDVAG FKRDGKSIFR ELALFQDYHG LPSFKKALVD FMSEIRGNKV TFDPNHIVLT AGSTSANETL 127
Q43166 WLTQNPDAAA FKRNGDSIFR DLALFQDYHG LPAFKDALVQ FMSEIRGNKV SFDSNKLVLT AGATSANETL 127
Conservation
Q42668 IFCLADPGDA FLIPSPYYAA FDRDLQWRTR AQIIPVHCNS SNNFQITREA LEVAYKKAEE SNIKVKGLII 207
Q43810 VFCLANPGDA FLLPSPYYPA NDRDLQWRTG AQIIPVHCNS SNGFKITREA LERSYAQAQE SNINVKGVLL 202
Q96579 AFCLADPGDA FLVPTPYYPG FDRDLRWRTG VQLFPVVCES CNDFKVTTKA LEEAYEKAQQ SNIKIKGLLI 207
Q43747 AFCLANPGDG FLVPTPYYPG FDRDLRWRTG VNLVPVTCHS SNGFKITAEA LDAAYENARV SNIPVKGLLI 207
Le-ACS4 ILCLADPGDA FLVPTPYYPG FNRDLRWRSG VQLLPISCKS CNNFKITIEA IEEAYEKGQQ ANVKIKGLIL 210
Q41688 MFCLAEQGDA FLLPTPYYPG FDRDLKWRTG AEIVPIQCTS SNNFQITEPA LKQAYEEARK RNLRVKGLLV 197
Q43166 MFCLADPGDA FLLPTPYYPG FDRDLKWRTG AEIVPIQCTS SNGFRITESA LEEAYKEAER RNLRVKGVLV 197
Conservation
Q42668 TNPSNPLGTI YDRDTLKSLV NFVNDNN-IH LICDEIYSAT VFKSPSFTSI AEIIEEMDH- -CKK-ELIHI 273
Q43810 TNPSNPLGTI LDRDTLKSIV SFVTDNN-IH LVIDEIYAAT VFVAPEFVSV SEILQEMDDT TCNP-DLIHI 270
Q96579 NNPSNPLGTL LDKDTLRDIV TFINSKN-IH LVCDEIYAAT VFDQPRFISV SEIVEDMIE- -CNK-DLIHI 273
Q43747 TNPSNPLGTT LDRDCLKSLV KFTNDKG-IH LIADEIYAAT TFGESEFISV AEVIDEIPD- -CNT-DLIHI 273
Le-ACS4 TNPCNPLGTI LDRDTLKKIS TFTNEHN-IH LVCDEIYAAT VFNSPKFVSI AEIINE--DN CINK-DLVHI 276
Q41688 TNPSNPLGTT MSRSELNLLV DFIKEKKDTH LISDEIYSGT VFSSPGFVSV MEILKERNDT -EEIWNRVHV 266
Q43166 TNPSNPLGTT LTKKELQLLL TFVSTKQ-IH LISDEIYSGT VFNSPKFVSV MEVLIENNYM YTEVWDRVHI 266
Conservation
Q42668 LSSLSKDMGL PGFRVGILYS YNDTVVSIAR KMSSFGLVSS QTQHLLAAML SDEEFVDNFL VENSKRLAKR 343
Q43810 VYSLSKDLGM PGFRVGIVYS FNDDVVSCAR KMSSFGLVST QTQHLLAAML SDDVFVERFL AER-RRLGRR 339
Q96579 VYSLSKDLGF PGFRVGIVYS YNDTVVNIAR KMSSFGLVSA QTQHLLASML SDEVFIDKFI AESSERLGER 343
Q43747 VYSLSKDMGL PGLRVGIVYS YNDRVVQIAR KMSSFGLVSS QTQHLIAKML SDEDFVDEFI RKSKLRLAER 343
Le-ACS4 VSSLSKDLGF PGFRVGIVYS FNDDVVNCAR KMSSFGLVST QTQHLLAFML SDDEFVEEFL IESAKRLRER 346
Q41688 VYSLSKDLGL PGFRVGAIYS ENDTVVAAAT KMSSFGLVSS QTQYLLSAML GDKKFARNYI VENQKRLKRR 336
Q43166 VYSLSKDLGL PGFRVGAIYS NDVMIVSAAT KMSSFGLISS QTQYLLSALL SDKKFTKKYV SENQKRLKKR 336
Conservation
Q42668 HARFTEELEK MGITCLNSNA GLFVWMDLRK LLKDQSFESE MELWRVIINE VKLNVSPGSS FHVTEPGWFR 413
Q43810 HGVFTKGLEE LGIGCLKSNA GLYFWMDLRK LLEEETFEAE MVLWKVIINE VKLNVSPGSS FHCVEPGWFR 409
Q96579 QGMFTKGLAE VGISTLKSNA GLFFWMDLRR LLKEATFDSE LELWRIIINE VKLNVSPGCS FHCSEPGWFR 413
Q43747 HAELTTGLDG LSIGWLKAGA GLFIWMDLRN LLKTATFDSE MELWRVIVHK VKLNVSPGGS CHCHEPGWFR 413
Le-ACS4 YEKFTRGLEE IGIKCLESNA GVYCWMDLRS LLKEATLDAE MSLWKLIINE VKLNVSPGSS FNCSEVGWFR 416
Q41688 QRMMVSGLQK AGISCLESNA GLFCWVDMRH LLHSNTFEAE MELWKKIVYQ VGLNISPGSS CHCTEPGWFR 406
Q43166 HEMLVGGLKQ IGIRCLESNA GLFCWVDMRH LLSSNTFDGE MELWKKIVYE VGLNISPGSS CHCTEPGWFR 406
Conservation
Q42668 VCFANMDDNT VGVALDRIHS FVGKIDKKEN S-TIPMPPKK KHRENKLRLS FSFSGRRYEE GNVLKSP-HM 481
Q43810 VCFANMDDET VHVALKRIRA FVRK---KEV G-----PVKR KRFMDNLNLR LSFSSLRYDE -SVMLSP-HI 469
Q96579 VCFANMDDET MRIALKRISY FV--LQPKGL N-NIA-AIKK QCSRRKLQIS LSFRRLDHE- --FMNSPAHS 476
Q43747 VCFANMDYQT METALERIRV FTSQTEEESL ILTKPMAKKK KCWQSSLRLS FK-DTRRFEE GFF--SP-HS 479
Le-ACS4 VCFANIDDQT MEIALARIRM FM----DAYN NVNKNGVMKN KHNGRGTTYD L--------- -----TP--- 465
Q41688 VCFANMSEET LTLAMKRLKN FVVESTCTQR SALFNGTSKR KSLTKWV-FR LSSRDHREQE ---------- 465
Q43166 ACFANMSEDT LNIAIQRLKA FVDSRDNKDD IQNQKHSNKK KSFSKWV-FR LSFNER--QR ---------- 463
Conservation
Q42668 M-SPHSPLVR AKT 493
Q43810 MVSTHSPLVR ART 482
Q96579 PMN--SPLVR --T 485
Q43747 PVPP-SPLVR AQI 491
Le-ACS4 --QMGSTMKM LLA 476
Q41688 ---------- -ER 467
Q43166 ---------- -ER 465
Conservation

Figure 2



ethylene production [nl h-1 g-1
12,00
10,00
8,00
6,00
4,00
2,00
0,00
2477 Pink
2477 Red
4043 Pink
4043 Red
4222 Pink
4222 Red
4303 Pink
4303 Red
5251 Pink
5251 Red
tapa Pink
tapa Red

Figure 3



% of fruits in red stage
100
90
80
70
60
50
40
30
20
10
0
0
5
10
15
20
25
30
35
Days [first fruit of Wt in breaker stage = Day 1]
Wild type
2477
4043
4222
4691
5251

Figure 4

```
                                                    Small domain

ACS4_WT_ID1      MDLETSEISN YKSSVVLSKL ASNEQHGENS PYFDGWKAYD NDPFHLVNNL 50
ACS4_2477_ID2    MDLETSEISN YKSSVVLSKL ASNEQHGENS PYFDGWKAYD NDPFHLVNNL
ACS4_4043_ID3    MDLETSEISN YKSSVVLSKL ASNEQHGENS PYFDGWKAYD NDPFHLVNNL
ACS4_4222_ID4    MDLETSEISN YKSSVVLSKL ASNEQHGENS PYFDGWKAYD NDPFHLVNNL
ACS4_4303_ID5    MDLETSEISN YKSSVVLSKL ASNEQHGENS PYFDGWKAYD NDPFHLVNNL
ACS4_4691_ID6    MDLETSEISN YKSSVVLSKL ASNEQHGENS PYFDGWKAYD NDPFHLVNNL
ACS4_5251_ID7    MDLETSEISN YKSSVVLSKL ASNEQHGENS PYFDGWKAYD NDPFHLVNNL

                                 Large domain

ACS4_WT_ID1      NGVIQMGLAE NQLSVDLIEE WIKRNPKASI CTNDGIESFR RIANFQDYHG 100
ACS4_2477_ID2    NGVIQMGLAE NQLSVDLIEE WIKRNPKASI CTNDGIESFR RIANFQDYHG
ACS4_4043_ID3    NGVIQMGLAE NQLSVDLIEE WIKRNPKASI CTNDGIESFR RIANFQDYHG
ACS4_4222_ID4    NGVIQMGLAE NQLSVDLIEE WIKRNPKASI CTNDGIESFR RIANFQDYHG
ACS4_4303_ID5    NGVIQMGLAE NQLSVDLIEE WIKRNPKASI CTNDGIESFR RIANFQDYHG
ACS4_4691_ID6    NGVIQMGLAE NQLSVDLIEE WIKRNPKASI CTNDGIESFR RIANFQDYHG
ACS4_5251_ID7    NGVIQMGLAE NQLSVDLIEE WIKRNPKASI CTNDGIESFR RIANFQDYHG


ACS4_WT_ID1      LPEFTNAIAK FMEKTRGGKV KFDAKRVVMA GGATGANETL ILCLADPGDA 150
ACS4_2477_ID2    LPEFTNAIAK FMEKTRGGKV KFDAKRVVMA GGATGANETL ILCLADPGDA
ACS4_4043_ID3    LPEFTNAIAK FMEKTRGGKV KFDAKRVVMA GGATGANETL ILCLADPGDA
ACS4_4222_ID4    LPEFTNAIAK FMEKTRGGKV KFDAKRVVMA GGATGANETL ILCLADPGDA
ACS4_4303_ID5    LPEFTNAIAK FMEKTRGGKV KFDAKRVVMA GGATGANETL ILCLADPGDA
ACS4_4691_ID6    LPEFTNAIAK FMEKTRGGKV KFDAKRVVMA GGATGANETL ILCLADPGDA
ACS4_5251_ID7    LPEFTNAIAK FMEKTRGGKV KFDAKRVVMA GGATGANETL ILCLADPGDA
```

```
ACS4_WT_ID1       FLVPTPYYPG FNRDLRWRSG VQLLPISCKS CNNFKITIEA IEEAYEKGQQ 200

ACS4_2477_ID2     FLVPTPYYPG FNRDLRWRSG VQLLPISCKS CNNFKITIEA IEEAYEKGQQ

ACS4_4043_ID3     FLVPTPYYPG FNRDLRWRSG VQLLPISCKS CNNFKITIEA IEEAYEKGQQ

ACS4_4222_ID4     FLVPTPYYPG FNRDLRWRSG VQLLPISCKS CNNFKITIEA IEEAYEKGQQ

ACS4_4303_ID5     FLVPTPYYPG FNRDLRWRSG VQLLPISCKS CNNFKITIEA IEEAYEKGQQ

ACS4_4691_ID6     FLVPTPYYPG FNRDLRWRSG VQLLPISCKS CNNFKITIEA IEEAYEKGQQ

ACS4_5251_ID7     FLVPTPYYPG FNRDLRWRSG VQLLPISCKS CNNFKITIEA IEEAYEKGQQ


ACS4_WT_ID1       ANVKIKGLIL TNPCNPLGTI LDRDTLKKIS TFTNEHNIHL VCDEIYAATV 250

ACS4_2477_ID2     ANVKIKGLIL TNPCNPLGTI LDRDTLKKIS TFTNEHNIHL VCDEIYAATV

ACS4_4043_ID3     ANVKIKGLIL TNPCNPLGTI LDRDTLKKIS TFTNEHNIHL VCDEIYAVTV

ACS4_4222_ID4     ANV*

ACS4_4303_ID5     ANVKIKGLIL TNPCNPLGTI LDRDTLKKIS TFTNEHNIHL VCDEIYAATV

ACS4_4691_ID6     ANVKIKGLIL TNPCNPLGTI LDRDTLKKIS TFTNEHNIHL VCDEIYAATE

ACS4_5251_ID7     ANVKIKGLIL TNPCNPLGTI LDRDTLKKIS TFTNEHNIHL VCDEIYAATV


ACS4_WT_ID1       FNSPKFVSIA EIINEDNCIN KDLVHIVSSL SKDLGFPGFR VGIVYSFNDD 300

ACS4_2477_ID2     FNSPKFVSIA EIINEDNCIN KDLVHIVSNL SKDLGFPGFR VGIVYSFNDD

ACS4_4043_ID3     FNSPKFVSIA EIINEDNCIN KDLVHIVSSL SKDLGFPGFR VGIVYSFNDD

ACS4_4222_ID4

ACS4_4303_ID5     FNSPKFVSIA EIINEDNCIN KDLVHIVSSL SKDLGFPGFR VGIVYSFNDD

ACS4_4691_ID6     FNSPKFVSIA EIINEDNCIN KDLVHIVSSL SKDLGFPGFR VGIVYSFNDD

ACS4_5251_ID7     FNSPKFVSIA EIINEDNCIN KDLVHIVSSL SKDLGFPGFR VGIVYSFNDD
```

Small domain

```
ACS4_WT_ID1      VVNCARKMSS FGLVSTQTQH LLAFMLSDDE FVEEFLIESA KRLRERYEKF 350
ACS4_2477_ID2    VVNCARKMSS FGLVSTQTQH LLAFMLSDDE FVEEFLIESA KRLRERYEKF
ACS4_4043_ID3    VVNCARKMSS FGLVSTQTQH LLAFMLSDDE FVEEFLIESA KRLRERYEKF
ACS4_4222_ID4
ACS4_4303_ID5    VVNCARKMSS FGLVSTQTQH FLAFMLSDDE FVEEFLIESA KRLRERYEKF
ACS4_4691_ID6    VVNCARKMSS FGLVSTQTQH LLAFMLSDDE FVEEFLIESA KRLRERYEKF
ACS4_5251_ID7    VVNCARKMSS FGLVSIQTQH LLAFMLSDDE FVEEFLIESA KRLRERYEKF

ACS4_WT_ID1      TRGLEEIGIK CLESNAGVYC WMDLRSLLKE ATLDAEMSLW KLIINEVKLN 400
ACS4_2477_ID2    TRGLEEIGIK CLESNAGVYC WMDLRSLLKE ATLDAEMSLW KLIINEVKLN
ACS4_4043_ID3    TRGLEEIGIK CLESNAGVYC WMDLRSLLKE ATLDAEMSLW KLIINEVKLN
ACS4_4222_ID4
ACS4_4303_ID5    TRGLEEIGIK CLESNAGVYC WMDLRSLLKE ATLDAEMSLW KLIINEVKLN
ACS4_4691_ID6    TRGLEEIGIK CLESNAGVYC WMDLRSLLKE ATLDAEMSLW KLIINEVKLN
ACS4_5251_ID7    TRGLEEIGIK CLESNAGVYC WMDLRSLLKE ATLDAEMSLW KLIINEVKLN

ACS4_WT_ID1      VSPGSSFNCS EVGWFRVCFA NIDDQTMEIA LARIRMFMDA YNNVNKNGVM 450
ACS4_2477_ID2    VSPGSSFNCS EVGWFRVCFA NIDDQTMEIA LARIRMFMDA YNNVNKNGVM
ACS4_4043_ID3    VSPGSSFNCS EVGWFRVCFA NIDDQTMEIA LARIRMFMDA YNNVNKNGVM
ACS4_4222_ID4
ACS4_4303_ID5    VSPGSSFNCS EVGWFRVCFA NIDDQTMEIA LARIRMFMDA YNNVNKNGVM
ACS4_4691_ID6    VSPGSSFNCS EVGWFRVCFA NIDDQTMEIA LARIRMFMDA YNNVNKNGVM
ACS4_5251_ID7    VSPGSSFNCS EVGWFRVCFA NIDDQTMEIA LARIRMFMDA YNNVNKNGVM
```

```
ACS4_WT_ID1      KNKHNGRGTT YDLTPQMGST MKMLLA

ACS4_2477_ID2    KNKHNGRGTT YDLTPQMGST MKMLLA

ACS4_4043_ID3    KNKHNGRGTT YDLTPQMGST MKMLLA

ACS4_4222_ID4

ACS4_4303_ID5    KNKHNGRGTT YDLTPQMGST MKMLLA

ACS4_4691_ID6    KNKHNGRGTT YDLTPQMGST MKMLLA

ACS4_5251_ID7    KNKHNGRGTT YDLTPQMGST MKMLLA
```

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 2005016504 A **[0013]**
- EP 1963505 A **[0091]**

Non-patent literature cited in the description

- **ALEXANDER.** *Journal of Experimental Botany,* 2002, vol. 53 (377), 2039-2055 **[0009]**
- **STEARNS ; GLICK.** *Biotechnology Advances,* 2003, vol. 21, 193-210 **[0010]**
- **CAPITANI et al.** a sequence alignment of eight ACS proteins. *Malus domestica, Phaseolus aureus, Solanum tuberosum, Pelargonium hortorum, Nicotiana tabacum, Cucumis melo, Lycopersicon esculentum, and Brassica oleracea* **[0011]**
- **CAPITANI et al.** *Journal of Molecular Biology,* 1999, vol. 294, 745-756 **[0011]**
- **BARRY et al.** *Plant Physiology,* 2000, vol. 123, 979-986 **[0012]**
- **YOKOTANI et al.** *Journal of Experimental Botany,* 2009, vol. 60, 3433-3442 **[0014]**
- **HENIKOFF ; HENIKOFF.** *PNAS,* 1992, vol. 89, 10915-10919 **[0042]**
- **CAPITANI et al.** *Journal of Molecular Biology,* 1999, vol. 194, 745-756 **[0043] [0080]**
- **HENIKOFF et al.** *Plant Physiology,* 2004, vol. 135, 630-636 **[0124]**
- **TILL et al.** *BMC Plant Biol,* 2007, vol. 7, 19 **[0124]**
- **TILL et al.** *Methods Mol Biol,* 2006, vol. 323, 127-35 **[0124]**
- **TILL et al.** *BMC Plant Biol,* 2004, vol. 4, 12 **[0124]**
- **TILL et al.** *Nat Protoc,* 2006, vol. 1, 2465-77 **[0124]**
- **COMAI et al.** *Plant J,* 2004, vol. 37, 778-86 **[0124]**
- **RIGOLA et al.** *PloS One,* March 2009, vol. 4 (3), e4761 **[0141]**